

## Chapter Nine

# SEQUENCES AND SERIES

## Contents

<b>9.1 Sequences . . . . .</b>	<b>464</b>
The Numerical Algebraic, and Graphical Viewpoint . . . . .	464
Defining Sequences Recursively . . . . .	465
Convergence of Sequences . . . . .	466
Convergence and Bounded Sequences . . . . .	466
<b>9.2 Geometric Series . . . . .</b>	<b>470</b>
Repeated Drug Dosage . . . . .	470
What Happens as $n \rightarrow \infty$ ? . . . .	471
The Geometric Series in General . . . . .	471
Sum of a Finite Geometric Series . . . . .	471
Sum of an Infinite Geometric Series . . . . .	472
Regular Deposits into a Savings Account . . . . .	473
What Happens as $n \rightarrow \infty$ ? . . . .	474
<b>9.3 Convergence of Series . . . . .</b>	<b>476</b>
Partial Sums and Convergence of Series . . . . .	476
Visualizing Series . . . . .	476
Comparison of Series and Integrals . . . . .	477
<b>9.4 Tests for Convergence . . . . .</b>	<b>481</b>
Comparison of Series . . . . .	481
Limit Comparison Test . . . . .	483
Series of Both Positive and Negative Terms . . . . .	484
Comparison with a Geometric Series: The Ratio Test . . . . .	485
Alternating Series . . . . .	486
Absolute and Conditional Convergence . . . . .	487
<b>9.5 Power Series and Interval of Convergence . . . . .</b>	<b>490</b>
Numerical and Graphical View of Convergence . . . . .	491
Intervals of Convergence . . . . .	492
What Happens at the Endpoints of the Interval of Convergence? . . . . .	495
<b>REVIEW PROBLEMS . . . . .</b>	<b>497</b>
<b>CHECK YOUR UNDERSTANDING . . . . .</b>	<b>501</b>
<b>PROJECTS . . . . .</b>	<b>502</b>

## 9.1 SEQUENCES

A sequence<sup>1</sup> is an infinite list of numbers  $s_1, s_2, s_3, \dots, s_n, \dots$ . We call  $s_1$  the first term,  $s_2$  the second term;  $s_n$  is the general term. For example, the sequence of squares,  $1, 4, 9, \dots, n^2, \dots$  can be denoted by the general term  $s_n = n^2$ . Thus, a sequence is a function whose domain is the positive integers, but it is traditional to denote the terms of a sequence using subscripts,  $s_n$ , rather than function notation,  $s(n)$ . In addition, we may talk about sequences whose general term has no simple formula, such as the sequence 3, 3.1, 3.14, 3.141, 3.1415,  $\dots$ , in which  $s_n$  gives the first  $n$  digits of  $\pi$ .

### The Numerical, Algebraic, and Graphical Viewpoint

Just as we can view a function algebraically, numerically, graphically, or verbally, we can view sequences in different ways. We may give an algebraic formula for the general term. We may give the numerical values of the first few terms of the sequence, suggesting a pattern for the later terms.

**Example 1** Give the first six terms of the following sequences:

(a)  $s_n = \frac{n(n+1)}{2}$

(b)  $s_n = \frac{n + (-1)^n}{n}$

**Solution**

(a) Substituting  $n = 1, 2, 3, 4, 5, 6$  into the formula for the general term, we get

$$\frac{1 \cdot 2}{2}, \frac{2 \cdot 3}{2}, \frac{3 \cdot 4}{2}, \frac{4 \cdot 5}{2}, \frac{5 \cdot 6}{2}, \frac{6 \cdot 7}{2} = 1, 3, 6, 10, 15, 21.$$

(b) Substituting  $n = 1, 2, 3, 4, 5, 6$  into the formula for the general term, we get

$$\frac{1-1}{1}, \frac{2+1}{2}, \frac{3-1}{3}, \frac{4+1}{4}, \frac{5-1}{5}, \frac{6+1}{6} = 0, \frac{3}{2}, \frac{2}{3}, \frac{5}{4}, \frac{4}{5}, \frac{7}{6}.$$

**Example 2** Give a general term for the following sequences:

(a) 1, 2, 4, 8, 16, 32,  $\dots$

(b)  $\frac{7}{2}, \frac{7}{5}, \frac{7}{8}, \frac{7}{11}, \frac{1}{2}, \frac{7}{17}, \dots$

**Solution**

Although the first six terms do not determine the sequence, we can sometimes use them to guess a possible formula for the general term.

(a) We have powers of 2, so we guess  $s_n = 2^n$ . When we check by substituting in  $n = 1, 2, 3, 4, 5, 6$ , we get 2, 4, 8, 16, 32, 64, instead of 1, 2, 4, 8, 16, 32. We fix our guess by subtracting 1 from the exponent, so the general term is

$$s_n = 2^{n-1}.$$

Substituting the first six values of  $n$  shows that the formula checks.

(b) In this sequence, the fifth term looks different from the others, whose numerators are all 7. We can fix this by rewriting  $1/2 = 7/14$ . The sequence of denominators is then 2, 5, 8, 11, 14, 17. This looks like a linear function with slope 3, so we expect the denominator has formula  $3n + k$  for some  $k$ . When  $n = 1$ , the denominator is 2, so

$$2 = 3 \cdot 1 + k \quad \text{giving} \quad k = -1$$

and the denominator of  $s_n$  is  $3n - 1$ . Our general term is then

$$s_n = \frac{7}{3n-1}.$$

To check this, evaluate  $s_n$  for  $n = 1, \dots, 6$ .

<sup>1</sup>In everyday English, the words “sequence” and “series” are used interchangeably. In mathematics, they have different meanings and cannot be interchanged.

and the denominator of  $s_n$  is  $3n - 1$ . Our general term is then

$$s_n = \frac{7}{3n - 1}.$$

To check this, evaluate  $s_n$  for  $n = 1, \dots, 6$ .

There are two ways to visualize a sequence. One is to plot points with  $n$  on the horizontal axis and  $s_n$  on the vertical axis. The other is to label points on a number line  $s_1, s_2, s_3, \dots$ . See Figure 9.1 for the sequence  $s_n = 1 + (-1)^n/n$ .




Figure 9.1: The sequence  $s_n = 1 + (-1)^n/n$

## Defining Sequences Recursively

Sequences can also be defined *recursively*, by giving an equation relating the  $n^{\text{th}}$  term to the previous terms and as many of the first terms as are needed to get started.

**Example 3** Give the first six terms of the recursively defined sequences.

- (a)  $s_n = s_{n-1} + 3$  for  $n > 1$  and  $s_1 = 4$
- (b)  $s_n = -3s_{n-1}$  for  $n > 1$  and  $s_1 = 2$
- (c)  $s_n = \frac{1}{2}(s_{n-1} + s_{n-2})$  for  $n > 2$  and  $s_1 = 0, s_2 = 1$
- (d)  $s_n = ns_{n-1}$  for  $n > 1$  and  $s_1 = 1$

**Solution** (a) When  $n = 2$ , we obtain  $s_2 = s_1 + 3 = 4 + 3 = 7$ . When  $n = 3$ , we obtain  $s_3 = s_2 + 3 = 7 + 3 = 10$ . In words, we obtain each term by adding 3 to the previous term. The first six terms are

$$4, 7, 10, 13, 16, 19.$$

(b) Each term is  $-3$  times the previous term, starting with  $s_1 = 2$ . We have  $s_2 = -3s_1 = -3 \cdot 2 = -6$  and  $s_3 = -3s_2 = -3(-6) = 18$ . Continuing, we get

$$2, -6, 18, -54, 162, -486.$$

(c) Each term is the average of the previous two terms, starting with  $s_1 = 0$  and  $s_2 = 1$ . We get  $s_3 = (s_2 + s_1)/2 = (1 + 0)/2 = 1/2$ . Then  $s_4 = (s_3 + s_2)/2 = ((1/2) + 1)/2 = 3/4$ . Continuing, we get

$$0, 1, \frac{1}{2}, \frac{3}{4}, \frac{5}{8}, \frac{11}{16}.$$

(d) Here  $s_2 = 2s_1 = 2 \cdot 1 = 2$  so  $s_3 = 3s_2 = 3 \cdot 2 = 6$  and  $s_4 = 4s_3 = 4 \cdot 6 = 24$ . Continuing gives

$$1, 2, 6, 24, 120, 720.$$

The general term of part (d) of the previous example is given by  $s_n = n(n-1)(n-2) \dots 3 \cdot 2 \cdot 1$ , which is denoted  $s_n = n!$  and is called  $n$  factorial.

We can also look at the first few terms of a sequence and try to guess a recursive definition by looking for a pattern.

**Example 4** Give a recursive definition of the following sequences.

- (a)  $1, 3, 7, 15, 31, 63, \dots$
- (b)  $1, 4, 9, 16, 25, 36, \dots$

**Solution** (a) Each term is twice the previous term plus one; for example  $7 = 2 \cdot 3 + 1$  and  $63 = 2 \cdot 31 + 1$ . Thus, a recursive definition is

$$s_n = 2s_{n-1} + 1 \text{ for } n > 1 \text{ and } s_1 = 1.$$

There are other ways to define the sequence recursively. We might notice, for example, that the differences of consecutive terms are powers of 2. Thus, we could also use

$$s_n = s_{n-1} + 2^{n-1} \quad \text{for } n > 1 \text{ and } s_1 = 1.$$

- (b) We recognize the terms as the squares of the positive integers, but we are looking for a recursive definition which relates consecutive terms. We see that

$$s_2 = s_1 + 3$$

$$s_3 = s_2 + 5$$

$$s_4 = s_3 + 7$$

$$s_5 = s_4 + 9,$$

so the differences between consecutive terms are consecutive odd integers. The difference between  $s_n$  and  $s_{n-1}$  is  $2n - 1$ , so a recursive definition is

$$s_n = s_{n-1} + 2n - 1, \text{ for } n > 1 \text{ and } s_1 = 1.$$

Recursively defined sequences, sometimes called recurrence relations, are powerful tools used frequently in computer science, as well as differential equations. Finding a formula for the general term can be surprisingly difficult.

## Convergence of Sequences

The limit of a sequence  $s_n$  as  $n \rightarrow \infty$  is defined the same way as the limit of a function  $f(x)$  as  $x \rightarrow \infty$ ; see also Problem 59.

The sequence  $s_1, s_2, s_3, \dots, s_n, \dots$  has a **limit**  $L$ , written  $\lim_{n \rightarrow \infty} s_n = L$ , if  $s_n$  is as close to  $L$  as we please whenever  $n$  is sufficiently large. If a limit,  $L$ , exists, we say the sequence **converges** to its limit  $L$ . If no limit exists, we say the sequence **diverges**.

To calculate the limit of a sequence, we use what we know about the limits of functions, including the properties in Theorem 1.2 and the following facts:

- The sequence  $s_n = x^n$  converges to 0 if  $|x| < 1$  and diverges if  $|x| > 1$
- The sequence  $s_n = 1/n^p$  converges to 0 if  $p > 0$

**Example 5** Do the following sequences converge or diverge? If a sequence converges, find its limit.

(a)  $s_n = (0.8)^n$                       (b)  $s_n = \frac{1 - e^{-n}}{1 + e^{-n}}$                       (c)  $s_n = 1 + (-1)^n$

**Solution**

- (a) Since  $0.8 < 1$ , the sequence converges by the first fact and the limit is 0.  
 (b) Since  $e^{-1} < 1$ , we have  $\lim_{n \rightarrow \infty} e^{-n} = \lim_{n \rightarrow \infty} (e^{-1})^n = 0$  by the first fact, so  $\lim_{n \rightarrow \infty} s_n = 1$ .  
 (c) Since  $(-1)^n$  alternates in sign, the sequence alternates between 0 and 2. Thus the sequence  $s_n$  diverges, since it does not get close to any fixed value.

## Convergence and Bounded Sequences

A sequence  $s_n$  is *bounded* if there are numbers  $K$  and  $M$  such that  $K \leq s_n \leq M$  for all terms. If  $\lim_{n \rightarrow \infty} s_n = L$ , then from some point on, the terms are bounded between  $L - 1$  and  $L + 1$ . Thus we have the following fact:

A convergent sequence is bounded.



On the other hand, a bounded sequence need not be convergent. In Example 5, we saw that  $1 + (-1)^n$  diverges, but it is bounded between 0 and 2. To ensure that a bounded sequence converges we need to rule out this sort of oscillation. The following theorem gives a condition that ensures convergence for a bounded sequence. A sequence  $s_n$  is called *monotone* if it is either increasing, that is  $s_n < s_{n+1}$  for all  $n$ , or decreasing, that is  $s_n > s_{n+1}$  for all  $n$ .

### Theorem 9.1: Convergence of a Monotone, Bounded Sequence

If a sequence  $s_n$  is bounded and monotone, it converges.

To understand this theorem graphically, see Figure 9.2. The sequence  $s_n$  is increasing and bounded above by  $M$ , so the values of  $s_n$  must “pile up” at some number less than or equal to  $M$ . This number is the limit.<sup>2</sup>




Figure 9.2: Values of  $s_n$  for  $n = 1, 2, \dots, 10$

As an example, the sequence  $s_n = (1 + 1/n)^n$  can be shown to be increasing and bounded (see Project 1 on page 502). Theorem 9.1 then guarantees that this sequence has a limit, which turns out to be  $e$ . (In fact, the sequence can be used to define  $e$ .)

**Example 6** If  $s_n = (1 + 1/n)^n$ , find  $s_{100}$  and  $s_{1000}$ . How many decimal places agree with  $e$ ?

**Solution** We have  $s_{100} = (1.01)^{100} = 2.7048$  and  $s_{1000} = (1.001)^{1000} = 2.7169$ . Since  $e = 2.7183\dots$ , we see that  $s_{100}$  agrees with  $e$  to one decimal place and  $s_{1000}$  agrees with  $e$  to two decimal places.

## Exercises and Problems for Section 9.1

### Exercises

In Exercises 1–6, find a formula for  $s_n$ ,  $n \geq 1$ .

1. 4, 8, 16, 32, 64, ...
2. 1, 3, 7, 15, 31, ...
3. 2, 5, 10, 17, 26, ...
4. 1, -3, 5, -7, 9, ...
5.  $1/3, 2/5, 3/7, 4/9, 5/11, \dots$
6.  $1/2, -1/4, 1/6, -1/8, 1/10, \dots$

For Exercises 7–12, find the first five terms of the sequence from the formula for  $s_n$ ,  $n \geq 1$ .

7.  $2^n + 1$
8.  $n + (-1)^n$
9.  $\frac{2n}{2n+1}$
10.  $(-1)^n \left(\frac{1}{2}\right)^n$
11.  $(-1)^{n+1} \left(\frac{1}{2}\right)^{n-1}$
12.  $\left(1 - \frac{1}{n+1}\right)^{n+1}$

### Problems


Do the sequences in Problems 13–24 converge or diverge? If a sequence converges, find its limit.

13.  $2^n$
14.  $(0.2)^n$
15.  $3 + e^{-2n}$
16.  $(-0.3)^n$
17.  $\frac{n}{10} + \frac{10}{n}$
18.  $\frac{2^n}{3^n}$
19.  $\frac{2n+1}{n}$
20.  $\frac{(-1)^n}{n}$
21.  $\frac{2^n}{n^3}$
22.  $\frac{2n + (-1)^n 5}{4n - (-1)^n 3}$
23.  $\frac{\sin n}{n}$
24.  $\cos(\pi n)$

<sup>2</sup>See the online supplement for a proof.

25. Match formulas (a)–(d) with graphs (I)–(IV).

- (a)  $s_n = 1 - 1/n$       (b)  $s_n = 1 + (-1)^n/n$   
 (c)  $s_n = 1/n$       (d)  $s_n = 1 + 1/n$




26. Match formulas (a)–(e) with descriptions (I)–(V) of the behavior of the sequence as  $n \rightarrow \infty$ .

- (a)  $s_n = n(n+1) - 1$   
 (b)  $s_n = 1/(n+1)$   
 (c)  $s_n = 1 - n^2$   
 (d)  $s_n = \cos(1/n)$   
 (e)  $s_n = (\sin n)/n$
- (I) Diverges to  $-\infty$   
 (II) Diverges to  $+\infty$   
 (III) Converges to 0 through positive numbers  
 (IV) Converges to 1  
 (V) Converges to 0 through positive and negative numbers

27. Match formulas (a)–(e) with graphs (I)–(V).

- (a)  $s_n = 2 - 1/n$   
 (b)  $s_n = (-1)^n 2 + 1/n$   
 (c)  $s_n = 2 + (-1)^n/n$   
 (d)  $s_n = 2 + 1/n$   
 (e)  $s_n = (-1)^n 2 + (-1)^n/n$



In Exercises 28–31, find the first six terms of the recursively defined sequence.

28.  $s_n = 2s_{n-1} + 3$  for  $n > 1$  and  $s_1 = 1$   
 29.  $s_n = s_{n-1} + n$  for  $n > 1$  and  $s_1 = 1$   
 30.  $s_n = s_{n-1} + \left(\frac{1}{2}\right)^{n-1}$  for  $n > 1$  and  $s_1 = 0$   
 31.  $s_n = s_{n-1} + 2s_{n-2}$  for  $n > 2$  and  $s_1 = 1, s_2 = 5$

In electrical engineering, a continuous function like  $f(t) = \sin t$ , where  $t$  is time in seconds, is referred to as an analog signal. To digitize the signal, we sample  $f(t)$  every  $\Delta t$  seconds to form the sequence  $s_n = f(n\Delta t)$ . For example, sampling  $f$  every  $1/10$  second produces the sequence  $\sin(1/10), \sin(2/10), \sin(3/10), \dots$ . In Problems 32–34, give the first 6 terms of a sampling of the signal every  $\Delta t$  seconds.

32.  $f(t) = (t-1)^2, \Delta t = 0.5$   
 33.  $f(t) = \cos 5t, \Delta t = 0.1$   
 34.  $f(t) = \frac{\sin t}{t}, \Delta t = 1$

To smooth a sequence,  $s_1, s_2, s_3, \dots$ , we replace each term  $s_n$  by  $t_n$ , the average of  $s_n$  with its neighboring terms

$$t_n = \frac{(s_{n-1} + s_n + s_{n+1})}{3} \text{ for } n > 1.$$

We start with  $t_1 = (s_1 + s_2)/2$ , since  $s_1$  has only one neighbor. For Problems 35–37, smooth the sequence once and then smooth the resulting sequence. What do you notice?

35. 18, -18, 18, -18, 18, -18, 18, ...  
 36. 0, 0, 0, 18, 0, 0, 0, 0, ...  
 37. 1, 2, 3, 4, 5, 6, 7, 8, ...

In Exercises 38–43, find a recursive definition for the sequence.

38. 1, 3, 5, 7, 9, ...      39. 2, 4, 6, 8, 10, ...  
 40. 3, 5, 9, 17, 33, ...      41. 1, 5, 14, 30, 55, ...  
 42. 1, 3, 6, 10, 15, ...      43.  $1, 2, \frac{3}{2}, \frac{5}{3}, \frac{8}{5}, \frac{13}{8}, \dots$

In Problems 44–46, show that the sequence  $s_n$  satisfies the recurrence relation.

44.  $s_n = 3n - 2$   
 $s_n = s_{n-1} + 3$  for  $n > 1$  and  $s_1 = 1$   
 45.  $s_n = n(n+1)/2$   
 $s_n = s_{n-1} + n$  for  $n > 1$  and  $s_1 = 1$   
 46.  $s_n = 2n^2 - n$   
 $s_n = s_{n-1} + 4n - 3$  for  $n > 1$  and  $s_1 = 1$

For a function  $f$ , define a sequence recursively by  $x_n = f(x_{n-1})$  for  $n > 1$  and  $x_1 = a$ . Depending on  $f$  and the starting value  $a$ , this sequence may converge to a limit  $L$ . If  $L$  exists, it has the property that  $f(L) = L$ . For the functions and starting values in Problems 47–50, use a calculator to see if the sequence converges. [To obtain the terms of the sequence, repeatedly push the function button.]

47.  $f(x) = \cos x, a = 0$       48.  $f(x) = e^{-x}, a = 0$

49.  $f(x) = \sin x, a = 1$       50.  $f(x) = \sqrt{x}, a = 0.5$

51. Let  $V_n$  be the number of new SUVs sold in the US in month  $n$ , where  $n = 1$  is January 2004. In terms of SUVs, what do the following represent?

- (a)  $V_{10}$   
 (b)  $V_n - V_{n-1}$   
 (c)  $\sum_{i=1}^{12} V_i$  and  $\sum_{i=1}^n V_i$

52. (a) Let  $s_n$  be the number of ancestors a person has  $n$  generations ago. What is  $s_1$ ?  $s_2$ ? Find a formula for  $s_n$ .

- (b) For which  $n$  is  $s_n$  greater than 6 billion, the current world population? What does this tell you about your ancestors?

53. For  $1 \leq n \leq 10$ , find a formula for  $p_n$ , the payment in year  $n$  on a loan of \$100,000. Interest is 5% per year, compounded annually, and payments are made at the end of each year for ten years. Each payment is \$10,000 plus the interest on the amount of money outstanding.

54. World oil consumption was 82.459 million barrels per day in 2005 and is increasing by about 1.3% per year.<sup>3</sup> Let  $c_n$  be daily world oil consumption  $n$  years after 2005.

- (a) Find a formula for  $c_n$ .  
 (b) Find and interpret  $c_n - c_{n-1}$ .  
 (c) What does the sum  $\sum_{n=1}^{18} 365c_n$  represent? (You do not need to compute this sum.)

55. (a) Cans are stacked in a triangle on a shelf. The bottom row contains  $k$  cans, the row above contains one can fewer, and so on, until the top row, which has one can. How many rows are there? Find  $a_n$ , the number of cans in the  $n^{\text{th}}$  row,  $1 \leq n \leq k$  (where the top row is  $n = 1$ ).

- (b) Let  $T_n$  be the total number of cans in the top  $n$  rows. Find a recurrence relation for  $T_n$  in terms of  $T_{n-1}$ .

- (c) Show that  $T_n = \frac{1}{2}n(n+1)$  satisfies the recurrence relation.

56. You are deciding whether to buy a new or a two-year-old car (of the same make) based on which will have cost you less when you resell it at the end of three years. Your cost consists of two parts: the loss in value of the car and the repairs. A new car costs \$20,000 and loses 12% of

its value each year. Repairs are \$400 the first year and increase by 18% each subsequent year.

- (a) For a new car, find the first three terms of the sequence  $d_n$  giving the depreciation (loss of value) in dollars in year  $n$ . Give a formula for  $d_n$ .

- (b) Find the first three terms of the sequence  $r_n$ , the repair cost in dollars for a new car in year  $n$ . Give a formula for  $r_n$ .

- (c) Find the total cost of owning a new car for three years.

- (d) Find the total cost of owning the two-year-old car for three years. Which should you buy?

57. The Fibonacci sequence first studied by the thirteenth century Italian mathematician Leonardo di Pisa, also known as Fibonacci, is defined recursively by

$$F_n = F_{n-1} + F_{n-2} \text{ for } n > 2 \text{ and } F_1 = 1, F_2 = 1.$$

The Fibonacci sequence occurs in many branches of mathematics and can be found in patterns of plant growth (phyllotaxis).

- (a) Find the first 12 terms.

- (b) Show that the sequence of successive ratios  $F_{n+1}/F_n$  appears to converge to a number  $r$  satisfying the equation  $r^2 = r + 1$ . (The number  $r$  was known as the golden ratio to the ancient Greeks.)

- (c) Let  $r$  satisfy  $r^2 = r + 1$ . Show that the sequence  $s_n = Ar^n$ , where  $A$  is constant, satisfies the Fibonacci equation  $s_n = s_{n-1} + s_{n-2}$  for  $n > 2$ .

58. This problem defines the Calkin-Wilf-Newman sequence of positive rational numbers. The sequence is remarkable because every positive rational number appears as one of its terms and none appears more than once. Every real number  $x$  can be written as an integer  $A$  plus a number  $B$  where  $0 \leq B < 1$ . For example, for  $x = 12/5 = 2 + 2/5$  we have  $A = 2$  and  $B = 2/5$ . For  $x = 3 = 3 + 0$  we have  $A = 3$  and  $B = 0$ . Define the function  $f(x)$  by

$$f(x) = A + (1 - B).$$

For example,  $f(12/5) = 2 + (1 - 2/5) = 13/5$  and  $f(3) = 3 + (1 - 0) = 4$ .

- (a) Evaluate  $f(x)$  for  $x = 25/8, 13/9$ , and  $\pi$ .

- (b) Find the first six terms of the recursively defined Calkin-Wilf-Newman sequence:  $s_n = 1/f(s_{n-1})$  for  $n > 1$  and  $s_1 = 1$ .

59. Write a definition for  $\lim_{n \rightarrow \infty} s_n = L$  similar to the  $\epsilon, \delta$  definition for  $\lim_{x \rightarrow a} f(x) = L$  in Section 1.8. Instead of  $\delta$ , you will need  $N$ , a value of  $n$ .

60. The sequence  $s_n$  is increasing, the sequence  $t_n$  converges, and  $s_n \leq t_n$  for all  $n$ . Show that  $s_n$  converges.

<sup>3</sup>www.bp.com, accessed May 14, 2007.

## 9.2 GEOMETRIC SERIES

This section introduces infinite series of constants, which are sums of the form

$$1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots$$

$$0.4 + 0.04 + 0.004 + 0.0004 + \cdots$$

The individual numbers,  $1, \frac{1}{2}, \frac{1}{3}, \dots$ , or  $0.4, 0.04, \dots$ , etc., are called *terms* in the series. To talk about the *sum* of the series, we must first explain how to add infinitely many numbers.

Let us look at the repeated administration of a drug. In this example, the terms in the series represent each dose; the sum of the series represents the drug level in the body in the long run.

### Repeated Drug Dosage

A person with an ear infection is told to take antibiotic tablets regularly for several days. Since the drug is being excreted by the body between doses, how can we calculate the quantity of the drug remaining in the body at any particular time?

To be specific, let's suppose the drug is ampicillin (a common antibiotic) taken in 250 mg doses four times a day (that is, every six hours). It is known that at the end of six hours, about 4% of the drug is still in the body. What quantity of the drug is in the body right after the tenth tablet? The fortieth?

Let  $Q_n$  represent the quantity, in milligrams, of ampicillin in the blood right after the  $n^{\text{th}}$  tablet. Then

$$Q_1 = 250 \qquad \qquad \qquad = 250 \text{ mg}$$

$$Q_2 = \underbrace{250(0.04)}_{\text{Remnants of first tablet}} + \underbrace{250}_{\text{New tablet}} \qquad \qquad \qquad = 260 \text{ mg}$$

$$Q_3 = Q_2(0.04) + 250 = (250(0.04) + 250)(0.04) + 250$$

$$= \underbrace{250(0.04)^2 + 250(0.04)}_{\text{Remnants of first and second tablets}} + \underbrace{250}_{\text{New tablet}} \qquad \qquad \qquad = 260.4 \text{ mg}$$

$$Q_4 = Q_3(0.04) + 250 = (250(0.04)^2 + 250(0.04) + 250)(0.04) + 250$$

$$= \underbrace{250(0.04)^3 + 250(0.04)^2 + 250(0.04)}_{\text{Remnants of first, second, and third tablets}} + \underbrace{250}_{\text{New tablet}} \qquad \qquad \qquad = 260.416 \text{ mg.}$$

Looking at the pattern that is emerging, we guess that

$$Q_5 = 250(0.04)^4 + 250(0.04)^3 + 250(0.04)^2 + 250(0.04) + 250$$

$$Q_{10} = 250(0.04)^9 + 250(0.04)^8 + \cdots + 250(0.04) + 250.$$

Notice that there are 10 terms in this sum—one for every tablet—but that the highest power of 0.04 is the ninth, because no tablet has been in the body for more than 9 six-hour time periods. (Do you see why?) Now suppose we actually want to find the numerical value of  $Q_{10}$ . It seems that we have to add 10 terms—and if we want the value of  $Q_{40}$ , we would be faced with adding 40 terms:

$$Q_{40} = 250(0.04)^{39} + 250(0.04)^{38} + \cdots + 250(0.04) + 250.$$

Fortunately, there's a better way. Let's start with  $Q_{10}$ .

$$Q_{10} = 250(0.04)^9 + 250(0.04)^8 + 250(0.04)^7 + \cdots + 250(0.04)^2 + 250(0.04) + 250.$$

Notice the remarkable fact that if you subtract  $(0.04)Q_{10}$  from  $Q_{10}$ , a great many terms (all but two, in fact) drop out. First multiplying by 0.04, we get

$$(0.04)Q_{10} = 250(0.04)^{10} + 250(0.04)^9 + 250(0.04)^8 + \cdots + 250(0.04)^3 + 250(0.04)^2 + 250(0.04).$$

Subtracting gives

$$Q_{10} - (0.04)Q_{10} = 250 - 250(0.04)^{10}.$$

Factoring  $Q_{10}$  on the left and solving for  $Q_{10}$  gives

$$\begin{aligned} Q_{10}(1 - 0.04) &= 250(1 - (0.04)^{10}) \\ Q_{10} &= \frac{250(1 - (0.04)^{10})}{1 - 0.04}. \end{aligned}$$

This is called the *closed-form* expression for  $Q_{10}$ . It is easy to evaluate on a calculator, giving  $Q_{10} = 260.42$  (to two decimal places). Similarly,  $Q_{40}$  is given in closed-form by

$$Q_{40} = \frac{250(1 - (0.04)^{40})}{1 - 0.04}.$$

Evaluating this on a calculator shows  $Q_{40} = 260.42$ , which is the same (to two decimal places) as  $Q_{10}$ . Thus after ten tablets, the value of  $Q_n$  appears to have stabilized at just over 260 mg.

Looking at the closed-forms for  $Q_{10}$  and  $Q_{40}$ , we can see that, in general,  $Q_n$  must be given by

$$Q_n = \frac{250(1 - (0.04)^n)}{1 - 0.04}.$$

### What Happens as $n \rightarrow \infty$ ?

What does this closed-form for  $Q_n$  predict about the long-run level of ampicillin in the body? As  $n \rightarrow \infty$ , the quantity  $(0.04)^n \rightarrow 0$ . In the long run, assuming that 250 mg continue to be taken every six hours, the level right after a tablet is taken is given by

$$Q_n = \frac{250(1 - (0.04)^n)}{1 - 0.04} \rightarrow \frac{250(1 - 0)}{1 - 0.04} = 260.42.$$

## The Geometric Series in General

In the previous example we encountered sums of the form  $a + ax + ax^2 + \cdots + ax^8 + ax^9$  (with  $a = 250$  and  $x = 0.04$ ). Such a sum is called a finite *geometric series*. A geometric series is one in which each term is a constant multiple of the one before. The first term is  $a$ , and the constant multiplier, or *common ratio* of successive terms, is  $x$ . (In our example,  $a = 250$  and  $x = 0.04$ .)

A **finite geometric series** has the form

$$a + ax + ax^2 + \cdots + ax^{n-2} + ax^{n-1}.$$

An **infinite geometric series** has the form

$$a + ax + ax^2 + \cdots + ax^{n-2} + ax^{n-1} + ax^n + \cdots.$$

The “ $\cdots$ ” at the end of the second series tells us that the series is going on forever—in other words, that it is infinite.

## Sum of a Finite Geometric Series

The same procedure that enabled us to find the closed-form for  $Q_{10}$  can be used to find the sum of any finite geometric series. Suppose we write  $S_n$  for the sum of the first  $n$  terms, which means up to the term containing  $x^{n-1}$ :

$$S_n = a + ax + ax^2 + \cdots + ax^{n-2} + ax^{n-1}.$$



Multiply  $S_n$  by  $x$ :

$$xS_n = ax + ax^2 + ax^3 + \cdots + ax^{n-1} + ax^n.$$

Now subtract  $xS_n$  from  $S_n$ , which cancels out all terms except for two, giving

$$\begin{aligned} S_n - xS_n &= a - ax^n \\ (1 - x)S_n &= a(1 - x^n). \end{aligned}$$

Provided  $x \neq 1$ , we can solve to find a closed form for  $S_n$  as follows:

The **sum of a finite geometric series** is given by

$$S_n = a + ax + ax^2 + \cdots + ax^{n-1} = \frac{a(1 - x^n)}{1 - x}, \quad \text{provided } x \neq 1.$$

Note that the value of  $n$  in the formula for  $S_n$  is the number of terms in the sum  $S_n$ .

## Sum of an Infinite Geometric Series

In the ampicillin example, we found the sum  $Q_n$  and then let  $n \rightarrow \infty$ . We do the same here. The sum  $Q_n$ , which shows the effect of the first  $n$  doses, is an example of a *partial sum*. The first three partial sums of the series  $a + ax + ax^2 + \cdots + ax^{n-1} + ax^n + \cdots$  are

$$\begin{aligned} S_1 &= a \\ S_2 &= a + ax \\ S_3 &= a + ax + ax^2. \end{aligned}$$

To find the sum,  $S$ , of this infinite series, we consider the partial sum,  $S_n$ , of the first  $n$  terms. The formula for the sum of a finite geometric series gives

$$S_n = a + ax + ax^2 + \cdots + ax^{n-1} = \frac{a(1 - x^n)}{1 - x}.$$

What happens to  $S_n$  as  $n \rightarrow \infty$ ? It depends on the value of  $x$ . If  $|x| < 1$ , then  $x^n \rightarrow 0$  as  $n \rightarrow \infty$ , so

$$\lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \frac{a(1 - x^n)}{1 - x} = \frac{a(1 - 0)}{1 - x} = \frac{a}{1 - x}.$$

Thus, provided  $|x| < 1$ , as  $n \rightarrow \infty$  the partial sums  $S_n$  approach a limit of  $a/(1 - x)$ . When this happens, we define the sum of the infinite geometric series to be that limit and say the series *converges* to  $a/(1 - x)$ .

For  $|x| < 1$ , the **sum of the infinite geometric series** is given by

$$S = a + ax + ax^2 + \cdots + ax^{n-1} + ax^n + \cdots = \frac{a}{1 - x}.$$

If, on the other hand,  $|x| > 1$ , then  $x^n$  and the partial sums have no limit as  $n \rightarrow \infty$  (if  $a \neq 0$ ). In this case, we say the series *diverges*. If  $x > 1$ , the terms in the series become larger and larger in magnitude, and the partial sums diverge to  $+\infty$  (if  $a > 0$ ) or  $-\infty$  (if  $a < 0$ ). When  $x < -1$ , the terms become larger in magnitude, the partial sums oscillate as  $n \rightarrow \infty$ , and the series diverges.

What happens when  $x = 1$ ? The series is

$$a + a + a + a + \cdots,$$

and if  $a \neq 0$ , the partial sums grow without bound, and the series does not converge. When  $x = -1$ , the series is

$$a - a + a - a + a - \cdots,$$

and, if  $a \neq 0$ , the partial sums oscillate between  $a$  and  $0$ , and the series does not converge.

**Example 1** For each of the following infinite geometric series, find several partial sums and the sum (if it exists).

$$(a) \quad 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \cdots \quad (b) \quad 1 + 2 + 4 + 8 + \cdots \quad (c) \quad 6 - 2 + \frac{2}{3} - \frac{2}{9} + \frac{2}{27} - \cdots$$

**Solution** (a) This series may be written

$$1 + \frac{1}{2} + \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^3 + \cdots$$

which we can identify as a geometric series with  $a = 1$  and  $x = \frac{1}{2}$ , so  $S = \frac{1}{1 - (1/2)} = 2$ .

Let's check this by finding the partial sums:

$$\begin{aligned} S_1 &= 1 \\ S_2 &= 1 + \frac{1}{2} = \frac{3}{2} = 2 - \frac{1}{2} \\ S_3 &= 1 + \frac{1}{2} + \frac{1}{4} = \frac{7}{4} = 2 - \frac{1}{4} \\ S_4 &= 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} = \frac{15}{8} = 2 - \frac{1}{8} \\ S_5 &= 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} = \frac{31}{16} = 2 - \frac{1}{16}. \end{aligned}$$

The formula for  $S_n$  gives

$$S_n = \frac{1 - (\frac{1}{2})^n}{1 - \frac{1}{2}} = 2 - \left(\frac{1}{2}\right)^{n-1}.$$

Thus, the partial sums are creeping up to the value  $S = 2$ , so  $S_n \rightarrow 2$  as  $n \rightarrow \infty$ .

(b) The partial sums of this geometric series (with  $a = 1$  and  $x = 2$ ) grow without bound, so the series has no sum:

$$\begin{aligned} S_1 &= 1 \\ S_2 &= 1 + 2 = 3 \\ S_3 &= 1 + 2 + 4 = 7 \\ S_4 &= 1 + 2 + 4 + 8 = 15 \\ S_5 &= 1 + 2 + 4 + 8 + 16 = 31. \end{aligned}$$

The formula for  $S_n$  gives

$$S_n = \frac{1 - 2^n}{1 - 2} = 2^n - 1.$$

(c) This is an infinite geometric series with  $a = 6$  and  $x = -\frac{1}{3}$ . The partial sums,

$$S_1 = 6.00, \quad S_2 = 4.00, \quad S_3 \approx 4.67, \quad S_4 \approx 4.44, \quad S_5 \approx 4.52, \quad S_6 \approx 4.49,$$

appear to be converging to 4.5. This turns out to be correct because the sum is

$$S = \frac{6}{1 - (-1/3)} = 4.5.$$

## Regular Deposits into a Savings Account

People who save money often do so by putting some fixed amount aside regularly. To be specific, suppose \$1000 is deposited every year in a savings account earning 5% a year, compounded annually. What is the balance,  $B_n$ , in dollars, in the savings account right after the  $n^{\text{th}}$  deposit?

As before, let's start by looking at the first few years:

$$B_1 = 1000$$

$$B_2 = B_1(1.05) + 1000 = \underbrace{1000(1.05)}_{\text{Original deposit}} + \underbrace{1000}_{\text{New deposit}}$$

$$B_3 = B_2(1.05) + 1000 = \underbrace{1000(1.05)^2 + 1000(1.05)}_{\text{First two deposits}} + \underbrace{1000}_{\text{New deposit}}$$

$$B_4 = B_3(1.05) + 1000 = \underbrace{1000(1.05)^3 + 1000(1.05)^2 + 1000(1.05)}_{\text{First three deposits}} + \underbrace{1000}_{\text{New deposit}}$$

Observing the pattern, we see

$$B_n = 1000(1.05)^{n-1} + 1000(1.05)^{n-2} + \cdots + 1000(1.05) + 1000.$$

So  $B_n$  is a finite geometric series with  $a = 1000$  and  $x = 1.05$ . Thus we have

$$B_n = \frac{1000(1 - (1.05)^n)}{1 - 1.05}.$$

We can rewrite this so that both the numerator and denominator of the fraction are positive:

$$B_n = \frac{1000((1.05)^n - 1)}{1.05 - 1}.$$

### What Happens as $n \rightarrow \infty$ ?

Common sense tells you that if you keep depositing \$1000 in an account and it keeps earning interest, your balance grows without bound. This is what the formula for  $B_n$  shows also:  $(1.05)^n \rightarrow \infty$  as  $n \rightarrow \infty$ , so  $B_n$  has no limit. (Alternatively, observe that the infinite geometric series of which  $B_n$  is a partial sum has  $x = 1.05$ , which is greater than 1, so the series does not converge.)

## Exercises and Problems for Section 9.2

### Exercises

In Exercises 1–10, decide which of the following are geometric series. For those which are, give the first term and the ratio between successive terms. For those which are not, explain why not.

- $5 - 10 + 20 - 40 + 80 - \cdots$
- $1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \cdots$
- $2 + 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \cdots$
- $1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \frac{1}{16} + \cdots$
- $1 + x + 2x^2 + 3x^3 + 4x^4 + \cdots$
- $1 + 2z + (2z)^2 + (2z)^3 + \cdots$
- $3 + 3z + 6z^2 + 9z^3 + 12z^4 + \cdots$
- $1 - x + x^2 - x^3 + x^4 - \cdots$
- $1 - y^2 + y^4 - y^6 + \cdots$
- $y^2 + y^3 + y^4 + y^5 + \cdots$

For each finite geometric series in Exercises 11–13, say how many terms are in the series and find its sum.

11.  $2 + 2(0.1) + 2(0.1)^2 + \cdots + 2(0.1)^{25}$

12.  $2(0.1) + 2(0.1)^2 + \cdots + 2(0.1)^{10}$

13.  $2(0.1)^5 + 2(0.1)^6 + \cdots + 2(0.1)^{13}$

Find the sum of the series in Exercises 14–17.

14.  $3 + \frac{3}{2} + \frac{3}{4} + \frac{3}{8} + \cdots + \frac{3}{2^{10}}$

15.  $-2 + 1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \frac{1}{16} - \cdots$

16.  $\sum_{n=4}^{\infty} \left(\frac{1}{3}\right)^n$

17.  $\sum_{n=4}^{20} \left(\frac{1}{3}\right)^n$

In Exercises 18–23, use the properties of geometric series to find the sum of the series. For what values of the variable does the series converge to this sum?

18.  $1 + z/2 + z^2/4 + z^3/8 + \cdots$

19.  $1 + 3x + 9x^2 + 27x^3 + \cdots$

20.  $y - y^2 + y^3 - y^4 + \cdots$

21.  $2 - 4z + 8z^2 - 16z^3 + \cdots$

22.  $3 + x + x^2 + x^3 + \cdots$

23.  $4 + y + y^2/3 + y^3/9 + \cdots$

## Problems

24. Bill invests \$200 at the start of each month for 24 months, starting now. If the investment yields 0.5% per month, compounded monthly, what is its value at the end of 24 months?
25. This problem shows another way of deriving the long-run ampicillin level. (See page 470.) In the long run the ampicillin levels off to  $Q$  mg right after each tablet is taken. Six hours later, right before the next dose, there will be less ampicillin in the body. However, if stability has been reached, the amount of ampicillin that has been excreted is exactly 250 mg because taking one more tablet raises the level back to  $Q$  mg. Use this to solve for  $Q$ .
26. On page 470, you saw how to compute the quantity  $Q_n$  mg of ampicillin in the body right after the  $n^{\text{th}}$  tablet of 250 mg, taken once every six hours.
- Do a similar calculation for  $P_n$ , the quantity of ampicillin (in mg) in the body right *before* the  $n^{\text{th}}$  tablet is taken.
  - Express  $P_n$  in closed form.
  - What is  $\lim_{n \rightarrow \infty} P_n$ ? Is this limit the same as  $\lim_{n \rightarrow \infty} Q_n$ ? Explain in practical terms why your answer makes sense.
27. Figure 9.3 shows the quantity of the drug atenolol in the blood as a function of time, with the first dose at time  $t = 0$ . Atenolol is taken in 50 mg doses once a day to lower blood pressure.




Figure 9.3

- If the half-life of atenolol in the blood is 6.3 hours, what percentage of the atenolol present at the start of a 24-hour period is still there at the end?
  - Find expressions for the quantities  $Q_0, Q_1, Q_2, Q_3, \dots$ , and  $Q_n$  shown in Figure 9.3. Write the expression for  $Q_n$  in closed-form.
  - Find expressions for the quantities  $P_1, P_2, P_3, \dots$ , and  $P_n$  shown in Figure 9.3. Write the expression for  $P_n$  in closed-form.
28. Draw a graph like that in Figure 9.3 for 250 mg of ampicillin taken every 6 hours, starting at time  $t = 0$ . Put on the graph the values of  $Q_1, Q_2, Q_3, \dots$  introduced in the text on page 470 and the values of  $P_1, P_2, P_3, \dots$  calculated in Problem 26.
29. A \$200,000 loan is to be repaid over 20 years in equal monthly installments of  $\$M$ , beginning at the end of the first month. Find the monthly payment if the loan is at an annual rate of 9%, compounded monthly. [Hint: Find an expression for the present value of the sum of all of the monthly payments, set it equal to \$200,000, and solve for  $M$ .]
30. (a) The total reserves of a non-renewable resource are 400 million tons. Annual consumption, currently 25 million tons per year, is expected to rise by 1% each year. After how many years will the reserves be exhausted?
- (b) Instead of increasing by 1% per year, suppose consumption was decreasing by a constant percentage per year. If existing reserves are never to be exhausted, what annual percentage reduction in consumption is required?
31. One way of valuing a company is to calculate the present value of all its future earnings. Suppose a farm expects to sell \$1000 worth of Christmas trees once a year forever, with the first sale in the immediate future. What is the present value of this Christmas tree business? Assume that the interest rate is 4% per year, compounded continuously.
32. Around January 1, 1993, Barbra Streisand signed a contract with Sony Corporation for \$2 million a year for 10 years. Suppose the first payment was made on the day of signing and that all other payments are made on the first day of the year. Suppose also that all payments are made into a bank account earning 4% a year, compounded annually.
- How much money was in the account
    - On the night of December 31, 1999?
    - On the day the last payment was made?
  - What was the present value of the contract on the day it was signed?
33. In theory, drugs that decay exponentially always leave a residue in the body. However, in practice, once the drug has been in the body for 5 half-lives, it is regarded as being eliminated.<sup>4</sup> If a patient takes a tablet of the same drug every 5 half-lives forever, what is the upper limit to the amount of drug that can be in the body?
34. This problem deals with the question of estimating the cumulative effect of a tax cut on a country's economy. Suppose the government proposes a tax cut totaling \$100 million. We assume that all the people who have extra money to spend would spend 80% of it and save 20%. Thus, of the extra income generated by the tax cut,  $\$100(0.8)$  million = \$80 million would be spent and so become extra income to someone else. Assume that these people also spend 80% of their additional income,

<sup>4</sup>[http://dr.pierce1.net/PDF/half\\_life.pdf](http://dr.pierce1.net/PDF/half_life.pdf), accessed on May 10, 2003.

or \$80(0.8) million, and so on. Calculate the total additional spending created by such a tax cut.

35. The government proposes a tax cut of \$100 million as in Problem 34, but that economists now predict that people will spend 90% of their extra income and save only 10%. How much additional spending would be generated by the tax cut under these assumptions?
36. A ball is dropped from a height of 10 feet and bounces. Each bounce is  $\frac{3}{4}$  of the height of the bounce before. Thus, after the ball hits the floor for the first time, the ball rises to a height of  $10(\frac{3}{4}) = 7.5$  feet, and after it hits the floor for the second time, it rises to a height of  $7.5(\frac{3}{4}) = 10(\frac{3}{4})^2 = 5.625$  feet. (Assume that there is no air resistance.)

- (a) Find an expression for the height to which the ball rises after it hits the floor for the  $n^{\text{th}}$  time.
- (b) Find an expression for the total vertical distance the

ball has traveled when it hits the floor for the first, second, third, and fourth times.

- (c) Find an expression for the total vertical distance the ball has traveled when it hits the floor for the  $n^{\text{th}}$  time. Express your answer in closed-form.
37. You might think that the ball in Problem 36 keeps bouncing forever since it takes infinitely many bounces. This is not true!
- (a) Show that a ball dropped from a height of  $h$  feet reaches the ground in  $\frac{1}{4}\sqrt{h}$  seconds. (Assume  $g = 32 \text{ ft/sec}^2$ )
- (b) Show that the ball in Problem 36 stops bouncing after

$$\frac{1}{4}\sqrt{10} + \frac{1}{2}\sqrt{10}\sqrt{\frac{3}{4}}\left(\frac{1}{1 - \sqrt{3/4}}\right) \approx 11 \text{ seconds.}$$

## 9.3 CONVERGENCE OF SERIES

We now consider general series in which each term  $a_n$  is a number. The series can be written compactly using a  $\sum$  sign as follows

$$\sum_{n=1}^{\infty} a_n = a_1 + a_2 + a_3 + \cdots + a_n + \cdots$$

For any particular values of  $a$  and  $x$ , the geometric series is such a series, with general term  $a_n = ax^{n-1}$ .

### Partial Sums and Convergence of Series

As in Section 9.2, we define the *partial sum*,  $S_n$ , of the first  $n$  terms of a series as

$$S_n = \sum_{i=1}^n a_i = a_1 + a_2 + \cdots + a_n.$$

To investigate the convergence of the series, we consider the sequence of partial sums

$$S_1, S_2, S_3, \dots, S_n, \dots$$

If  $S_n$  has a limit as  $n \rightarrow \infty$ , then we define the sum of the series to be that limit.

If the sequence  $S_n$  of partial sums converges to  $S$ , so  $\lim_{n \rightarrow \infty} S_n = S$ , then we say the series  $\sum_{n=1}^{\infty} a_n$  **converges** and that its sum is  $S$ . We write  $\sum_{n=1}^{\infty} a_n = S$ . If  $\lim_{n \rightarrow \infty} S_n$  does not exist, we say that the series **diverges**.

### Visualizing Series

We can visualize the terms of a series as in Figure 9.4. In this figure, we assume  $a_n \geq 0$  for all  $n$ , so each rectangle has area  $a_n$ . Then the series converges if the total area of the rectangles is finite and the sum of the series is the total area of the rectangles. This is similar to an improper integral  $\int_0^{\infty} f(x) dx$ , in which the area under the graph of  $f$  can be finite, even on an infinite interval.






Figure 9.4: Height and area of the  $n^{\text{th}}$  rectangle is  $a_n$

Here are some properties that are useful in determining whether or not a series converges.

### Theorem 9.2: Convergence Properties of Series

1. If  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  converge and if  $k$  is a constant, then
  - $\sum_{n=1}^{\infty} (a_n + b_n)$  converges to  $\sum_{n=1}^{\infty} a_n + \sum_{n=1}^{\infty} b_n$ .
  - $\sum_{n=1}^{\infty} k a_n$  converges to  $k \sum_{n=1}^{\infty} a_n$ .
2. Changing a finite number of terms in a series does not change whether or not it converges, although it may change the value of its sum if it does converge.
3. If  $\lim_{n \rightarrow \infty} a_n \neq 0$  or  $\lim_{n \rightarrow \infty} a_n$  does not exist, then  $\sum_{n=1}^{\infty} a_n$  diverges.
4. If  $\sum_{n=1}^{\infty} a_n$  diverges, then  $\sum_{n=1}^{\infty} k a_n$  diverges if  $k \neq 0$ .

For proofs of these properties, see Problems 39–42. As for improper integrals, the convergence of a series is determined by its behavior for large  $n$ . (See the “behaves like” principle on page 381.) From Property 2 we see that, if  $N$  is a positive integer, then  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=N}^{\infty} a_n$  either both converge or both diverge. Thus, if all we care about is the convergence of a series, we can omit the limits and write  $\sum a_n$ .

**Example 1** Does the series  $\sum (1 - e^{-n})$  converge?

**Solution** Since the terms in the series,  $a_n = 1 - e^{-n}$  tend to 1, not 0, as  $n \rightarrow \infty$ , the series diverges by Property 3 of Theorem 9.2.

## Comparison of Series and Integrals

We investigate the convergence of some series by comparison with an improper integral. The *harmonic series* is the infinite series

$$\sum_{n=1}^{\infty} \frac{1}{n} = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots + \frac{1}{n} + \cdots$$

Convergence of this sum would mean that the sequence of partial sums

$$S_1 = 1, \quad S_2 = 1 + \frac{1}{2}, \quad S_3 = 1 + \frac{1}{2} + \frac{1}{3}, \quad \cdots, \quad S_n = 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n}, \quad \cdots$$

tends to a limit as  $n \rightarrow \infty$ . Let's look at some values:

$$S_1 = 1, \quad S_{10} \approx 2.93, \quad S_{100} \approx 5.19, \quad S_{1000} \approx 7.49, \quad S_{10000} \approx 9.79.$$

The growth of these partial sums is slow, but they do in fact grow without bound, so the harmonic series diverges. This is justified in the following example and in Problem 46.

**Example 2** Show that the harmonic series  $1 + 1/2 + 1/3 + 1/4 + \dots$  diverges.

**Solution** The idea is to approximate  $\int_1^\infty (1/x) dx$  by a left-hand sum, where the terms  $1, 1/2, 1/3, \dots$  are heights of rectangles of base 1. In Figure 9.5, the sum of the areas of the 3 rectangles is larger than the area under the curve between  $x = 1$  and  $x = 4$ , and the same kind of relationship holds for the first  $n$  rectangles. Thus, we have

$$S_n = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} > \int_1^{n+1} \frac{1}{x} dx = \ln(n+1).$$

Since  $\ln(n+1)$  gets arbitrarily large as  $n \rightarrow \infty$ , so do the partial sums,  $S_n$ . Thus, the partial sums have no limit, so the series diverges.




Figure 9.5: Comparing the harmonic series to  $\int_1^\infty (1/x) dx$

Notice that the harmonic series diverges, even though  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} (1/n) = 0$ .

**Example 3** By comparison with the improper integral  $\int_1^\infty (1/x^2) dx$ , show that the following series converges:

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = 1 + \frac{1}{4} + \frac{1}{9} + \dots$$

**Solution** Since we want to show that  $\sum_{n=1}^{\infty} 1/n^2$  converges, we want to show that the partial sums of this series tend to a limit. We do this by showing that the sequence of partial sums increases and is bounded above, so Theorem 9.1 applies.

Each successive partial sum is obtained from the previous one by adding one more term in the series. Since all the terms are positive, the sequence of partial sums is increasing.

To show that the partial sums of  $\sum_{n=1}^{\infty} 1/n^2$  are bounded, we consider the right-hand sum represented by the area of the rectangles in Figure 9.6. We start at  $x = 1$ , since the area under the curve is infinite for  $0 \leq x \leq 1$ . The shaded rectangles in Figure 9.6 suggest that:

$$\frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \dots + \frac{1}{n^2} \leq \int_1^\infty \frac{1}{x^2} dx.$$

The area under the graph is finite, since

$$\int_1^\infty \frac{1}{x^2} dx = \lim_{b \rightarrow \infty} \int_1^b \frac{1}{x^2} dx = \lim_{b \rightarrow \infty} \left( -\frac{1}{b} + 1 \right) = 1.$$




Figure 9.6: Comparing  $\sum_{n=1}^{\infty} 1/n^2$  to  $\int_1^{\infty} (1/x^2) dx$

To get  $S_n$ , we add 1 to both sides, giving

$$S_n = 1 + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \cdots + \frac{1}{n^2} \leq 1 + \int_1^{\infty} \frac{1}{x^2} dx = 2.$$

Thus, the sequence of partial sums is bounded above by 2. Hence, by Theorem 9.1 the sequence of partial sums converges, so the series converges.

Notice that we have shown that the series in the Example 3 converges, but we have not found its sum. The integral gives us a bound on the partial sums, but it does not give us the limit of the partial sums. Euler proved the remarkable fact that the sum is  $\pi^2/6$ .

The method of Examples 2 and 3 can be used to prove the following theorem. See Problem 45.

### Theorem 9.3: The Integral Test

Suppose  $a_n = f(n)$ , where  $f(x)$  is decreasing and positive.

- If  $\int_1^{\infty} f(x) dx$  converges, then  $\sum a_n$  converges.
- If  $\int_1^{\infty} f(x) dx$  diverges, then  $\sum a_n$  diverges.

Suppose  $f(x)$  is continuous. Then if  $f(x)$  is positive and decreasing for all  $x$  beyond some point, say  $c$ , the integral test can be used.

The integral test allows us to analyze a family of series, the  $p$ -series, and see how convergence depends on the parameter  $p$ .

**Example 4** For what values of  $p$  does the series  $\sum_{n=1}^{\infty} 1/n^p$  converge?

**Solution** If  $p \leq 0$ , the terms in the series  $a_n = 1/n^p$  do not tend to 0 as  $n \rightarrow \infty$ . Thus the series diverges for  $p \leq 0$ .

If  $p > 0$ , we compare  $\sum_{n=1}^{\infty} 1/n^p$  to the integral  $\int_1^{\infty} 1/x^p dx$ . In Example 3 of Section 7.7 we saw that the integral converges if  $p > 1$  and diverges if  $p \leq 1$ . By the integral test, we conclude that  $\sum 1/n^p$  converges if  $p > 1$  and diverges if  $p \leq 1$ .

We can summarize Example 4 as follows:

The  $p$ -series  $\sum_{n=1}^{\infty} 1/n^p$  converges if  $p > 1$  and diverges if  $p \leq 1$ .

## Exercises and Problems for Section 9.3

## Exercises

In Exercises 1–3, find the first five terms of the sequence of partial sums.

$$1. \sum_{n=1}^{\infty} n \quad 2. \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \quad 3. \sum_{n=1}^{\infty} \frac{1}{n(n+1)}$$

Use the integral test to decide whether the series in Exercises 4–7 converge or diverge.

$$4. \sum_{n=1}^{\infty} \frac{1}{n^3} \quad 5. \sum_{n=1}^{\infty} \frac{n}{n^2 + 1} \\ 6. \sum_{n=1}^{\infty} \frac{1}{e^n} \quad 7. \sum_{n=2}^{\infty} \frac{1}{n(\ln n)^2}$$

8. Use comparison with  $\int_1^{\infty} x^{-3} dx$  to show that  $\sum_{n=2}^{\infty} 1/n^3$  converges to a number less than or equal to  $1/2$ .

9. Use comparison with  $\int_0^{\infty} 1/(x^2 + 1) dx$  to show that  $\sum_{n=1}^{\infty} 1/(n^2 + 1)$  converges to a number less than or equal to  $\pi/2$ .

Explain why the integral test cannot be used to decide if the series in Exercises 10–12 converge or diverge.

$$10. \sum_{n=1}^{\infty} n^2 \quad 11. \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \quad 12. \sum_{n=1}^{\infty} e^{-n} \sin n$$

## Problems

Do the series in Problems 13–32 converge or diverge?

$$13. \sum_{n=0}^{\infty} \frac{3}{n+2} \quad 14. \sum_{n=0}^{\infty} \frac{4}{2n+1} \\ 15. \sum_{n=0}^{\infty} \frac{2}{\sqrt{2+n}} \quad 16. \sum_{n=0}^{\infty} \frac{2n}{1+n^4} \\ 17. \sum_{n=0}^{\infty} \frac{2n}{(1+n^2)^2} \quad 18. \sum_{n=0}^{\infty} \frac{2n}{\sqrt{4+n^2}} \\ 19. \sum_{n=1}^{\infty} \frac{3}{(2n-1)^2} \quad 20. \sum_{n=1}^{\infty} \frac{4}{(2n+1)^3} \\ 21. \sum_{n=0}^{\infty} \frac{3}{n^2+4} \quad 22. \sum_{n=0}^{\infty} \frac{2}{1+4n^2} \\ 23. \sum_{n=1}^{\infty} \frac{n}{n+1} \quad 24. \sum_{n=0}^{\infty} \frac{n+1}{2n+3} \\ 25. \sum_{n=1}^{\infty} \left(\frac{1}{2}\right)^n + \left(\frac{2}{3}\right)^n \quad 26. \sum_{n=1}^{\infty} \left(\left(\frac{3}{4}\right)^n + \frac{1}{n}\right) \\ 27. \sum_{n=1}^{\infty} \frac{n+2^n}{n2^n} \quad 28. \sum_{n=1}^{\infty} \frac{\ln n}{n} \\ 29. \sum_{n=1}^{\infty} \frac{1}{n(1+\ln n)} \quad 30. \sum_{n=3}^{\infty} \frac{n+1}{n^2+2n+2} \\ 31. \sum_{n=0}^{\infty} \frac{1}{n^2+2n+2} \quad 32. \sum_{n=2}^{\infty} \frac{n \ln n + 4}{n^2}$$

33. Show that  $\sum_{n=1}^{\infty} \frac{1}{\ln(2^n)}$  diverges.

34. Show that  $\sum_{n=1}^{\infty} \frac{1}{(\ln(2^n))^2}$  converges.

35. (a) Find the partial sum,  $S_n$ , of  $\sum_{n=1}^{\infty} \ln\left(\frac{n+1}{n}\right)$ .

(b) Does the series in part (a) converge or diverge?

36. (a) Show  $r^{\ln n} = n^{\ln r}$  for positive numbers  $n$  and  $r$ .

(b) For what values  $r > 0$  does  $\sum_{n=1}^{\infty} r^{\ln n}$  converge?

37. Consider the series  $\sum_{k=1}^{\infty} \frac{1}{k(k+1)} = \frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \cdots$ .

(a) Show that  $\frac{1}{k} - \frac{1}{k+1} = \frac{1}{k(k+1)}$ .

(b) Use part (a) to find the partial sums  $S_3$ ,  $S_{10}$ , and  $S_n$ .

(c) Use part (b) to show that the sequence of partial sums  $S_n$ , and therefore the series, converges to 1.

38. Consider the series

$$\sum_{k=1}^{\infty} \ln\left(\frac{k(k+2)}{(k+1)^2}\right) = \ln\left(\frac{1 \cdot 3}{2 \cdot 2}\right) + \ln\left(\frac{2 \cdot 4}{3 \cdot 3}\right) + \cdots$$

(a) Show that the partial sum of the first three nonzero terms  $S_3 = \ln(5/8)$ .

(b) Show that the partial sum  $S_n = \ln\left(\frac{n+2}{2(n+1)}\right)$ .

(c) Use part (b) to show that the partial sums  $S_n$ , and therefore the series, converge to  $\ln(1/2)$ .

39. Show that if  $\sum a_n$  and  $\sum b_n$  converge and if  $k$  is a constant, then  $\sum(a_n + b_n)$ ,  $\sum(a_n - b_n)$ , and  $\sum ka_n$  converge.

40. Let  $N$  be a positive integer. Show that if  $a_n = b_n$  for  $n \geq N$ , then  $\sum a_n$  and  $\sum b_n$  either both converge, or both diverge.

41. Show that if  $\sum a_n$  converges, then  $\lim_{n \rightarrow \infty} a_n = 0$ . [Hint: Consider  $\lim_{n \rightarrow \infty} (S_n - S_{n-1})$ , where  $S_n$  is the  $n^{\text{th}}$  partial sum.]

42. Show that if  $\sum a_n$  diverges and  $k \neq 0$ , then  $\sum ka_n$  diverges.
43. The series  $\sum a_n$  converges. Explain, by looking at partial sums, why the series  $\sum (a_{n+1} - a_n)$  also converges.
44. The series  $\sum a_n$  diverges. Give examples that show the series  $\sum (a_{n+1} - a_n)$  could converge or diverge.
45. In this problem, you will justify the integral test. Suppose  $c \geq 0$  and  $f(x)$  is a decreasing positive function, defined for all  $x \geq c$ , with  $f(n) = a_n$  for all  $n$ .
- (a) Suppose  $\int_c^\infty f(x) dx$  converges. By considering rectangles under the graph of  $f$ , show that  $\sum a_n$  converges. [Hint: See Example 3 on page 478.]
- (b) Suppose that  $\int_c^\infty f(x) dx$  diverges. By considering rectangles above the graph of  $f$ , show that  $\sum a_n$  diverges. [Hint: See Example 2 on page 478.]
46. Consider the following grouping of terms in the harmonic series:
- $$1 + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right) + \left(\frac{1}{9} + \frac{1}{10} + \cdots + \frac{1}{16}\right) + \cdots$$
- (a) Show that the sum of each group of fractions is more than  $1/2$ .
- (b) Explain why this shows that the harmonic series does not converge.
47. Show that  $\sum_{n=2}^\infty \frac{1}{n \ln n}$  diverges.
- (a) Using the integral test.
- (b) By considering the grouping of terms
- $$\left(\frac{1}{2 \ln 2}\right) + \left(\frac{1}{3 \ln 3} + \frac{1}{4 \ln 4}\right) + \left(\frac{1}{5 \ln 5} + \frac{1}{6 \ln 6} + \frac{1}{7 \ln 7} + \frac{1}{8 \ln 8}\right) + \cdots$$
48. Consider the sequence given by
- $$a_n = \left(1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n}\right) - \ln n.$$
- (a) Show that  $a_n$  is positive for all  $n$ . [Use a left-sum approximation to  $\int_1^n (1/x) dx$  with  $\Delta x = 1$ .]
- (b) Show that  $a_{n+1} < a_n$  for all  $n$ . [Use one rectangle to give a right approximation to  $\int_n^{n+1} (1/x) dx$ .]
- (c) Explain why  $\lim_{n \rightarrow \infty} a_n$  exists.
- (d) The number  $\gamma = \lim_{n \rightarrow \infty} a_n$  is called *Euler's constant*. Estimate  $\gamma$  to two decimal places by computing  $a_{200}$ .
49. On page 479, we gave Euler's result
- $$\sum_{n=1}^\infty \frac{1}{n^2} = \frac{\pi^2}{6}.$$
- (a) Find the sum of the first 20 terms of this series. Give your answer to three decimal places.
- (b) Use your answer to estimate  $\pi$ . Give your answer to two decimal places.
- (c) Repeat parts (a) and (b) with 100 terms.
- (d) Use a right sum approximation to bound the error in approximating  $\pi^2/6$  by  $\sum_{n=1}^{20} (1/n^2)$  and by  $\sum_{n=1}^{100} (1/n^2)$ .
50. This problem approximates  $e$  using
- $$e = \sum_{n=0}^\infty \frac{1}{n!}.$$
- (a) Find a lower bound for  $e$  by evaluating the first five terms of the series.
- (b) Show that  $1/n! \leq 1/2^{n-1}$  for  $n \geq 1$ .
- (c) Find an upper bound for  $e$  using part (b).
51. In this problem we investigate how fast the partial sums  $S_N = 1^5 + 2^5 + 3^5 + \cdots + N^5$  of the divergent series  $\sum_{n=1}^\infty n^5$  grow as  $N$  gets larger and larger. Show that
- (a)  $S_N > N^6/6$  by considering the right-hand Riemann sum for  $\int_0^N x^5 dx$  with  $\Delta x = 1$ .
- (b)  $S_N < ((N+1)^6 - 1)/6$  by considering the left-hand Riemann sum for  $\int_1^{N+1} x^5 dx$  with  $\Delta x = 1$ .
- (c)  $\lim_{N \rightarrow \infty} S_N/(N^6/6) = 1$ . We say that  $S_N$  is asymptotic to  $N^6/6$  as  $N$  goes to infinity.

## 9.4 TESTS FOR CONVERGENCE

### Comparison of Series

In Section 7.8, we compared two integrals to decide whether an improper integral converged. In Theorem 9.3 we compared an integral and a series. Now we compare two series.

#### Theorem 9.4: Comparison Test

Suppose  $0 \leq a_n \leq b_n$  for all  $n$  beyond a certain value.

- If  $\sum b_n$  converges, then  $\sum a_n$  converges.
- If  $\sum a_n$  diverges, then  $\sum b_n$  diverges.



Since  $a_n \leq b_n$ , the plot of the  $a_n$  lies under the plot of the  $b_n$ . (See Figure 9.7.) The comparison test says that if the total area for  $\sum b_n$  is finite, then the total area for  $\sum a_n$  is finite also. If the total area for  $\sum a_n$  is not finite, then neither is the total area for  $\sum b_n$ .




Figure 9.7: Each  $a_n$  is represented by the area of a dark rectangle, and each  $b_n$  by a dark plus a light rectangle

**Example 1** Use the comparison test to determine whether the series  $\sum_{n=1}^{\infty} \frac{1}{n^3 + 1}$  converges.

**Solution** For  $n \geq 1$ , we know that  $n^3 \leq n^3 + 1$ , so

$$0 \leq \frac{1}{n^3 + 1} \leq \frac{1}{n^3}.$$

Thus, every term in the series  $\sum_{n=1}^{\infty} 1/(n^3 + 1)$  is less than or equal to the corresponding term in  $\sum_{n=1}^{\infty} 1/n^3$ . Since we saw that  $\sum_{n=1}^{\infty} 1/n^3$  converges as a  $p$ -series with  $p > 1$ , we know that  $\sum_{n=1}^{\infty} 1/(n^3 + 1)$  converges.

**Example 2** Decide whether the following series converge: (a)  $\sum_{n=1}^{\infty} \frac{n-1}{n^3+3}$  (b)  $\sum_{n=1}^{\infty} \frac{6n^2+1}{2n^3-1}$ .

**Solution** (a) Since the convergence is determined by the behavior of the terms for large  $n$ , we observe that

$$\frac{n-1}{n^3+3} \text{ behaves like } \frac{n}{n^3} = \frac{1}{n^2} \text{ as } n \rightarrow \infty.$$

Since  $\sum 1/n^2$  converges, we guess that  $\sum (n-1)/(n^3+3)$  converges. To confirm this, we use the comparison test. Since a fraction increases if its numerator is made larger or its denominator is made smaller, we have

$$0 \leq \frac{n-1}{n^3+3} \leq \frac{n}{n^3} = \frac{1}{n^2} \text{ for all } n \geq 1.$$

Thus, the series  $\sum (n-1)/(n^3+3)$  converges by comparison with  $\sum 1/n^2$ .

(b) First, we observe that

$$\frac{6n^2+1}{2n^3-1} \text{ behaves like } \frac{6n^2}{2n^3} = \frac{3}{n} \text{ as } n \rightarrow \infty.$$

Since  $\sum 1/n$  diverges, so does  $\sum 3/n$ , and we guess that  $\sum (6n^2 + 1)/(2n^3 - 1)$  diverges. To confirm this, we use the comparison test. Since a fraction decreases if its numerator is made smaller or its denominator is made larger, we have

$$0 \leq \frac{6n^2}{2n^3} \leq \frac{6n^2 + 1}{2n^3 - 1},$$

so

$$0 \leq \frac{3}{n} \leq \frac{6n^2 + 1}{2n^3 - 1}.$$

Thus, the series  $\sum (6n^2 + 1)/(2n^3 - 1)$  diverges by comparison with  $\sum 3/n$ .

### Limit Comparison Test

The convergence or divergence of a series  $\sum a_n$  is determined by the values of  $a_n$  as  $n \rightarrow \infty$ . This often enables us to predict convergence or divergence by looking at the long-run behavior of  $a_n$ .

**Example 3** Predict convergence or divergence of

$$\sum \frac{n^2 - 5}{n^3 + n + 2}.$$

**Solution** As  $n \rightarrow \infty$ , the highest power terms in the numerator and denominator,  $n^2$  and  $n^3$ , dominate. Thus the term

$$a_n = \frac{n^2 - 5}{n^3 + n + 2}$$

behaves, as  $n \rightarrow \infty$ , like

$$\frac{n^2}{n^3} = \frac{1}{n}.$$

Since the harmonic series  $\sum 1/n$  diverges, we guess that  $\sum a_n$  also diverges.

The comparison test is needed to confirm a prediction of convergence or divergence. Because the comparison test requires showing that  $a_n \leq b_n$ , we may prefer to use the following test, which avoids these inequalities.

#### Theorem 9.5: Limit Comparison Test

Suppose  $a_n > 0$  and  $b_n > 0$  for all  $n$ . If

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c \quad \text{where } c > 0,$$

then the two series  $\sum a_n$  and  $\sum b_n$  either both converge or both diverge.

The limit  $\lim_{n \rightarrow \infty} a_n/b_n = c$  captures the idea that  $a_n$  “behaves like”  $cb_n$  as  $n \rightarrow \infty$ .

**Example 4** Use the limit comparison test to determine if the following series converge or diverge.

(a)  $\sum \frac{n^2 + 6}{n^4 - 2n + 3}$

(b)  $\sum \sin\left(\frac{1}{n}\right)$

**Solution** (a) We take  $a_n = \frac{n^2 + 6}{n^4 - 2n + 3}$ . Because  $a_n$  behaves like  $\frac{n^2}{n^4} = \frac{1}{n^2}$  as  $n \rightarrow \infty$  we take  $b_n = 1/n^2$ . We have

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{n^4 + 6n^2}{n^4 - 2n + 3} = 1.$$

The limit comparison test applies with  $c = 1$ . Since  $\sum 1/n^2$  converges, the limit comparison test shows that  $\sum \frac{n^2 + 6}{n^4 - 2n + 3}$  also converges.

(b) Since  $\sin(x) \approx x$  for  $x$  near 0, we know that  $\sin(\frac{1}{n})$  behaves like  $1/n$  as  $n \rightarrow \infty$ . We apply the limit comparison test with  $a_n = \sin(\frac{1}{n})$  and  $b_n = 1/n$ . We have

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\sin(1/n)}{1/n} = 1.$$

Thus  $c = 1$  and since  $\sum 1/n$  diverges, the series  $\sum \sin(1/n)$  also diverges.

## Series of Both Positive and Negative Terms

If  $\sum a_n$  has both positive and negative terms, then its plot has rectangles lying both above and below the horizontal axis. See Figure 9.8. The total area of the rectangles is no longer equal to  $\sum a_n$ . However, it is still true that if the total area of the rectangles above and below the axis is finite, then the series converges. The area of the  $n^{\text{th}}$  rectangle is  $|a_n|$ , so we have:

### Theorem 9.6: Convergence of Absolute Values Implies Convergence

If  $\sum |a_n|$  converges, then so does  $\sum a_n$ .

Problem 87 shows how to prove this result.




Figure 9.8: Representing a series with positive and negative terms

**Example 5** Explain how we know that the following series converges

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2} = 1 - \frac{1}{4} + \frac{1}{9} - \cdots$$

**Solution** Writing  $a_n = (-1)^{n-1}/n^2$ , we have

$$|a_n| = \left| \frac{(-1)^{n-1}}{n^2} \right| = \frac{1}{n^2}.$$

The  $p$ -series  $\sum 1/n^2$  converges, since  $p > 1$ , so  $\sum (-1)^{n-1}/n^2$  converges.

## Comparison with a Geometric Series: The Ratio Test

A geometric series  $\sum a_n$  has the property that the ratio  $a_{n+1}/a_n$  is constant for all  $n$ . For many other series, this ratio, although not constant, tends to a constant as  $n$  increases. In some ways, such series behave like geometric series. In particular, a geometric series converges if the ratio  $|a_{n+1}/a_n| < 1$ . A non-geometric series also converges if the ratio  $|a_{n+1}/a_n|$  tends to a limit which is less than 1. This idea leads to the following test.

### Theorem 9.7: The Ratio Test

For a series  $\sum a_n$ , suppose the sequence of ratios  $|a_{n+1}|/|a_n|$  has a limit:

$$\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = L.$$

- If  $L < 1$ , then  $\sum a_n$  converges.
- If  $L > 1$ , or if  $L$  is infinite,<sup>5</sup> then  $\sum a_n$  diverges.
- If  $L = 1$ , the test does not tell us anything about the convergence of  $\sum a_n$ .

**Proof** Here are the main steps in the proof. Suppose  $\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = L < 1$ . Let  $x$  be a number between  $L$  and 1. Then for all sufficiently large  $n$ , say for all  $n \geq k$ , we have

$$\frac{|a_{n+1}|}{|a_n|} < x.$$

Then,

$$\begin{aligned} |a_{k+1}| &< |a_k|x, \\ |a_{k+2}| &< |a_{k+1}|x < |a_k|x^2, \\ |a_{k+3}| &< |a_{k+2}|x < |a_k|x^3, \end{aligned}$$

and so on. Thus, writing  $a = |a_k|$ , we have for  $i = 1, 2, 3, \dots$ ,

$$|a_{k+i}| < ax^i.$$

Now we can use the comparison test:  $\sum |a_{k+i}|$  converges by comparison with the geometric series  $\sum ax^i$ . Since  $\sum |a_{k+i}|$  converges, Theorem 9.6 tells us that  $\sum a_{k+i}$  converges. So, by property 2 of Theorem 9.2, we see that  $\sum a_n$  converges too.

If  $L > 1$ , then for sufficiently large  $n$ , say  $n \geq m$ ,

$$|a_{n+1}| > |a_n|,$$

so the sequence  $|a_m|, |a_{m+1}|, |a_{m+2}|, \dots$  is increasing. Thus,  $\lim_{n \rightarrow \infty} a_n \neq 0$ , so  $\sum a_n$  diverges (by Theorem 9.2, property 3). The argument in the case that  $|a_{n+1}|/|a_n|$  is unbounded is similar.

**Example 6** Show that the following series converges:<sup>6</sup>

$$\sum_{n=1}^{\infty} \frac{1}{n!} = 1 + \frac{1}{2!} + \frac{1}{3!} + \dots$$

**Solution** Since  $a_n = 1/n!$  and  $a_{n+1} = 1/(n+1)!$ , we have

$$\frac{|a_{n+1}|}{|a_n|} = \frac{1/(n+1)!}{1/n!} = \frac{n!}{(n+1)!} = \frac{n(n-1)(n-2) \cdots 2 \cdot 1}{(n+1)n(n-1) \cdots 2 \cdot 1}.$$

<sup>5</sup>That is, the sequence  $|a_{n+1}|/|a_n|$  grows without bound.

<sup>6</sup>We define  $2!$  to be  $2 \cdot 1 = 2$ . Similarly,  $3! = 3 \cdot 2 \cdot 1 = 6$  and  $n! = n \cdot (n-1) \cdots 2 \cdot 1$ . We also define  $0! = 1$ .

We cancel  $n(n-1)(n-2) \cdots 2 \cdot 1$ , giving

$$\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = \lim_{n \rightarrow \infty} \frac{n!}{(n+1)!} = \lim_{n \rightarrow \infty} \frac{1}{n+1} = 0.$$

Because the limit is 0, which is less than 1, the ratio test tells us that  $\sum_{n=1}^{\infty} 1/n!$  converges. In Chapter 10, we see that the sum is  $e - 1$ .

**Example 7** What does the ratio test tell us about the convergence of the following two series?

$$\sum_{n=1}^{\infty} \frac{1}{n} \quad \text{and} \quad \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}.$$

**Solution** Because  $|(-1)^n| = 1$ , in both cases we have  $\lim_{n \rightarrow \infty} |a_{n+1}/a_n| = \lim_{n \rightarrow \infty} n/(n+1) = 1$ . Thus, the ratio test does not tell us anything about the convergence of either series. In fact, the first series is the harmonic series, which diverges. Example 8 will show that the second series converges.

## Alternating Series

A series is called an *alternating series* if the terms alternate in sign. For example,

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots + \frac{(-1)^{n-1}}{n} + \cdots.$$

The convergence of an alternating series can often be determined using the following test:

### Theorem 9.8: Alternating Series Test

A series of the form

$$\sum_{n=1}^{\infty} (-1)^{n-1} a_n = a_1 - a_2 + a_3 - a_4 + \cdots + (-1)^{n-1} a_n + \cdots$$

converges if

$$0 < a_{n+1} < a_n \quad \text{for all } n \quad \text{and} \quad \lim_{n \rightarrow \infty} a_n = 0.$$

Although we do not prove this result, we can see why it is reasonable. The first partial sum,  $S_1 = a_1$ , is positive. The second,  $S_2 = a_1 - a_2$ , is still positive, since  $a_2 < a_1$ , but  $S_2$  is smaller than  $S_1$ . (See Figure 9.9.) The next sum,  $S_3 = a_1 - a_2 + a_3$ , is greater than  $S_2$  but smaller than  $S_1$ . The partial sums oscillate back and forth, and since the distance between them tends to 0, they eventually converge.




Figure 9.9: Partial sums,  $S_1, S_2, S_3, S_4$  of an alternating series

**Example 8** Show that the following alternating harmonic series converges:

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}.$$



Solution

We have  $a_n = 1/n$  and  $a_{n+1} = 1/(n+1)$ . Thus,

$$a_{n+1} = \frac{1}{n+1} < \frac{1}{n} = a_n \quad \text{for all } n, \quad \text{and} \quad \lim_{n \rightarrow \infty} 1/n = 0.$$

Thus, the hypothesis of Theorem 9.8 is satisfied, so the alternating harmonic series converges.

Suppose  $S$  is the sum of an alternating series, so  $S = \lim_{n \rightarrow \infty} S_n$ . Then  $S$  is trapped between any two consecutive partial sums, say  $S_3$  and  $S_4$  or  $S_4$  and  $S_5$  so

$$S_2 < S_4 < \cdots < S < \cdots < S_3 < S_1.$$

Thus, the error in using  $S_n$  to approximate the true sum  $S$  is less than the distance from  $S_n$  to  $S_{n+1}$ , which is  $a_{n+1}$ . Stated symbolically, we have the following result:

### Theorem 9.9: Error Bounds for Alternating Series

Let  $S_n = \sum_{i=1}^n (-1)^{i-1} a_i$  be the  $n^{\text{th}}$  partial sum of an alternating series and let  $S = \lim_{n \rightarrow \infty} S_n$ .

Suppose that  $0 < a_{n+1} < a_n$  for all  $n$  and  $\lim_{n \rightarrow \infty} a_n = 0$ . Then

$$|S - S_n| < a_{n+1}.$$

Thus, provided  $S_n$  converges to  $S$  by the alternating series test, the error in using  $S_n$  to approximate  $S$  is less than the magnitude of the first term of the series which is omitted in the approximation.

#### Example 9

Estimate the error in approximating the sum of the alternating harmonic series  $\sum_{n=1}^{\infty} (-1)^{n-1}/n$  by the sum of the first nine terms.

Solution

The ninth partial sum is given by

$$S_9 = 1 - \frac{1}{2} + \frac{1}{3} - \cdots + \frac{1}{9} = 0.7456 \dots$$

The first term omitted is  $-1/10$ , with magnitude 0.1. By Theorem 9.9, we know that the true value of the sum differs from  $0.7456 \dots$  by less than 0.1.

## Absolute and Conditional Convergence

We say that the series  $\sum a_n$  is

- **absolutely convergent** if  $\sum a_n$  and  $\sum |a_n|$  both converge.
- **conditionally convergent** if  $\sum a_n$  converges but  $\sum |a_n|$  diverges.

Conditionally convergent series rely on cancellation between positive and negative terms for their convergence.

*Example:* The series  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2}$  is absolutely convergent because the series converges and the  $p$ -series  $\sum 1/n^2$  also converges.

*Example:* The series  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}$  is conditionally convergent because the series converges but the harmonic series  $\sum 1/n$  diverges.

## Exercises and Problems for Section 9.4

## Exercises

Use the comparison test to confirm the statements in Exercises 1–3.

- $\sum_{n=4}^{\infty} \frac{1}{n}$  diverges, so  $\sum_{n=4}^{\infty} \frac{1}{n-3}$  diverges.
- $\sum_{n=1}^{\infty} \frac{1}{n^2}$  converges, so  $\sum_{n=1}^{\infty} \frac{1}{n^2+2}$  converges.
- $\sum_{n=1}^{\infty} \frac{1}{n^2}$  converges, so  $\sum_{n=1}^{\infty} \frac{e^{-n}}{n^2}$  converges.

In Exercises 4–7, use end behavior to compare the series to a  $p$ -series and predict whether the series converges or diverges.

- $\sum_{n=1}^{\infty} \frac{n^3+1}{n^4+2n^3+2n}$
- $\sum_{n=1}^{\infty} \frac{n+4}{n^3+5n-3}$
- $\sum_{n=1}^{\infty} \frac{1}{n^4+3n^3+7}$
- $\sum_{n=1}^{\infty} \frac{n-4}{\sqrt{n^3+n^2+8}}$

Use the comparison test to determine whether the series in Exercises 8–13 converge.

- $\sum_{n=1}^{\infty} \frac{1}{3^n+1}$
- $\sum_{n=1}^{\infty} \frac{1}{n^4+e^n}$
- $\sum_{n=2}^{\infty} \frac{1}{\ln n}$
- $\sum_{n=1}^{\infty} \frac{n^2}{n^4+1}$
- $\sum_{n=1}^{\infty} \frac{n \sin^2 n}{n^3+1}$
- $\sum_{n=1}^{\infty} \frac{2^n+1}{n2^n-1}$

Use the ratio test to decide if the series in Exercises 14–19 converge or diverge.

- $\sum_{n=1}^{\infty} \frac{1}{(2n)!}$
- $\sum_{n=1}^{\infty} \frac{(n!)^2}{(2n)!}$
- $\sum_{n=1}^{\infty} \frac{(2n)!}{n!(n+1)!}$
- $\sum_{n=1}^{\infty} \frac{1}{r^n n!}, r > 0$
- $\sum_{n=1}^{\infty} \frac{1}{ne^n}$
- $\sum_{n=0}^{\infty} \frac{2^n}{n^3+1}$

Which of the the series in Exercises 20–23 are alternating?

- $\sum_{n=1}^{\infty} (-1)^n \left(2 - \frac{1}{n}\right)$
- $\sum_{n=1}^{\infty} \cos(n\pi)$

- $\sum_{n=1}^{\infty} (-1)^n \cos(n\pi)$
- $\sum_{n=1}^{\infty} (-1)^n \cos n$

Use the alternating series test to show that the series in Exercises 24–27 converge.

- $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{\sqrt{n}}$
- $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{2n+1}$
- $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2+2n+1}$
- $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{e^n}$

In Exercises 28–34, determine whether the series is absolutely convergent, conditionally convergent, or divergent.

- $\sum \frac{(-1)^n}{2^n}$
- $\sum \frac{(-1)^n}{2n}$
- $\sum (-1)^n \left(1 + \frac{1}{n^2}\right)$
- $\sum \frac{(-1)^n}{n^4+7}$
- $\sum \frac{(-1)^{n-1}}{n \ln n}$
- $\sum (-1)^{n-1} \arcsin\left(\frac{1}{n}\right)$
- $\sum \frac{(-1)^{n-1} \arctan(1/n)}{n^2}$

In Exercises 35–43, use the limit comparison test to determine whether the series converges or diverges.

- $\sum_{n=1}^{\infty} \frac{5n+1}{3n^2}$ , by comparing to  $\sum_{n=1}^{\infty} \frac{1}{n}$
- $\sum_{n=1}^{\infty} \left(\frac{1+n}{3n}\right)^n$ , by comparing to  $\sum_{n=1}^{\infty} \left(\frac{1}{3}\right)^n$   
[Hint:  $\lim_{n \rightarrow \infty} (1 + 1/n)^n = e$ .]
- $\sum \left(1 - \cos \frac{1}{n}\right)$ , by comparing to  $\sum 1/n^2$
- $\sum \frac{1}{n^4-7}$
- $\sum \frac{n+1}{n^2+2}$
- $\sum \frac{n^3-2n^2+n+1}{n^4-2}$
- $\sum \frac{2^n}{3^n-1}$
- $\sum \frac{1}{2\sqrt{n}+\sqrt{n+2}}$
- $\sum \left(\frac{1}{2n-1} - \frac{1}{2n}\right)$

## Problems

Explain why the comparison test cannot be used to decide if the series in Problems 44–45 converge or diverge.

$$44. \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2}$$

$$45. \sum_{n=1}^{\infty} \sin n$$

Explain why the ratio test cannot be used to decide if the series in Problems 46–47 converge or diverge.

$$46. \sum_{n=1}^{\infty} (-1)^n$$

$$47. \sum_{n=1}^{\infty} \sin n$$

Explain why the alternating series test cannot be used to decide if the series in Problems 48–50 converge or diverge.

$$48. \sum_{n=1}^{\infty} (-1)^{n-1} n$$

$$49. \sum_{n=1}^{\infty} (-1)^{n-1} \sin n$$

$$50. \sum_{n=1}^{\infty} (-1)^{n-1} \left(2 - \frac{1}{n}\right)$$

Use a computer or calculator to investigate the behavior of the partial sums of the alternating series in Problems 51–53. Which ones converge? Confirm convergence using the alternating series test. If a series converges, estimate its sum.

$$51. 1 - 2 + 3 - 4 + 5 + \cdots + (-1)^n (n+1) + \cdots$$

$$52. 1 - 0.1 + 0.01 - 0.001 + \cdots + (-1)^n 10^{-n} + \cdots$$

$$53. 1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \cdots + (-1)^n \frac{1}{n!} + \cdots$$

Determine which of the series in Problems 54–72 converge.

$$54. \sum_{n=1}^{\infty} \frac{8^n}{n!}$$

$$55. \sum_{n=1}^{\infty} \frac{n2^n}{3^n}$$

$$56. \sum_{n=0}^{\infty} \frac{(0.1)^n}{n!}$$

$$57. \sum_{n=1}^{\infty} \frac{(n-1)!}{n^2}$$

$$58. \sum_{n=0}^{\infty} e^{-n}$$

$$59. \sum_{n=1}^{\infty} e^n$$

$$60. \sum_{n=1}^{\infty} \frac{(2n)!}{(n!)^2}$$

$$61. \sum_{n=1}^{\infty} \frac{1}{n^2} \tan\left(\frac{1}{n}\right)$$

$$62. \sum_{n=1}^{\infty} \frac{n+1}{n^3+6}$$

$$63. \sum_{n=1}^{\infty} \frac{5n+2}{2n^2+3n+7}$$

$$64. \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{\sqrt{3n-1}}$$

$$65. \sum_{n=1}^{\infty} \frac{(-1)^{n-1} 2^n}{n^2}$$

$$66. \sum_{n=1}^{\infty} \frac{\sin n}{n^2}$$

$$67. \sum_{n=1}^{\infty} \frac{\sin n^2}{n^2}$$

$$68. \sum_{n=1}^{\infty} \frac{\cos(n\pi)}{n}$$

$$69. \sum_{n=2}^{\infty} \frac{n+2}{n^2-1}$$

$$70. \sum_{n=2}^{\infty} \frac{3}{\ln n^2}$$

$$71. \sum_{n=1}^{\infty} \frac{1}{\sqrt{n^2(n+2)}}$$

$$72. \sum_{n=1}^{\infty} \frac{n(n+1)}{\sqrt{n^3+2n^2}}$$

For what values of  $a$  do the series in Problems 73–77 converge?

$$73. \sum_{n=1}^{\infty} \left(\frac{2}{n}\right)^a$$

$$74. \sum_{n=1}^{\infty} \left(\frac{2}{a}\right)^n, a > 0$$

$$75. \sum_{n=1}^{\infty} (\ln a)^n, a > 0$$

$$76. \sum_{n=1}^{\infty} \frac{\ln n}{n^a}$$

$$77. \sum_{n=1}^{\infty} (-1)^n \arctan\left(\frac{a}{n}\right), a > 0$$

The series in Problems 78–80 converge by the alternating series test. Use Theorem 9.9 to find how many terms give a partial sum,  $S_n$ , within 0.01 of the sum,  $S$ , of the series.

$$78. \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}$$

$$79. \sum_{n=1}^{\infty} \left(-\frac{2}{3}\right)^{n-1}$$

$$80. \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{(2n)!}$$

81. Suppose  $0 \leq b_n \leq 2^n \leq a_n$  and  $0 \leq c_n \leq 2^{-n} \leq d_n$  for all  $n$ . Which of the series  $\sum a_n$ ,  $\sum b_n$ ,  $\sum c_n$ , and  $\sum d_n$  definitely converge, which definitely diverge?

82. Given two convergent series  $\sum a_n$  and  $\sum b_n$ , we know that the term-by-term sum  $\sum (a_n + b_n)$  converges. What about the series formed by taking the products of the terms  $\sum a_n \cdot b_n$ ? This problem shows that it may or may not converge.

(a) Show that if  $\sum a_n = \sum 1/n^2$  and  $\sum b_n = \sum 1/n^3$ , then  $\sum a_n \cdot b_n$  converges.

(b) Explain why  $\sum (-1)^n / \sqrt{n}$  converges.

(c) Show that if  $a_n = b_n = (-1)^n / \sqrt{n}$ , then  $\sum a_n \cdot b_n$  does not converge.

83. Suppose that  $b_n > 0$  and  $\sum b_n$  converges. Show that if  $\lim_{n \rightarrow \infty} a_n/b_n = 0$  then  $\sum a_n$  converges.
84. Suppose that  $b_n > 0$  and  $\sum b_n$  diverges. Show that if  $\lim_{n \rightarrow \infty} a_n/b_n = \infty$  then  $\sum a_n$  diverges.
85. A series  $\sum a_n$  of positive terms (that is,  $a_n > 0$ ) can be used to form another series  $\sum b_n$  where each term  $b_n$  is the average of the first  $n$  terms of the original series, that is,  $b_n = (a_1 + a_2 + \cdots + a_n)/n$ . Show that  $\sum b_n$  does not converge (even if  $\sum a_n$  does). [Hint: Compare  $\sum b_n$  to a multiple of the harmonic series.]
86. Show that if  $\sum |a_n|$  converges, then  $\sum (-1)^n a_n$  converges.
87. (a) For a series  $\sum a_n$ , show that  $0 \leq a_n + |a_n| \leq 2|a_n|$ .  
 (b) Use part (a) to show that if  $\sum |a_n|$  converges, then  $\sum a_n$  converges.

Problems 88–89 introduce the *root test* for convergence. Given a series  $\sum a_n$  of positive terms (that is,  $a_n > 0$ ) such that the root  $\sqrt[n]{a_n}$  has a limit  $r$  as  $n \rightarrow \infty$ ,

- if  $r < 1$ , then  $\sum a_n$  converges
- if  $r > 1$ , then  $\sum a_n$  diverges
- if  $r = 1$ , then  $\sum a_n$  could converge or diverge.

(This test works since  $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = r$  tells us that the series is comparable to a geometric series with ratio  $r$ .) Use this test to determine the behavior of the series.

$$88. \sum_{n=1}^{\infty} \left(\frac{2}{n}\right)^n$$

$$89. \sum_{n=1}^{\infty} \left(\frac{5n+1}{3n^2}\right)^n$$

## 9.5 POWER SERIES AND INTERVAL OF CONVERGENCE

In Section 9.2 we saw that the geometric series  $\sum ax^n$  converges for  $-1 < x < 1$  and diverges otherwise. This section studies the convergence of more general series constructed from powers. Chapter 10 shows how such power series are used to approximate functions such as  $e^x$ ,  $\sin x$ ,  $\cos x$ , and  $\ln x$ .

A **power series** about  $x = a$  is a sum of constants times powers of  $(x - a)$ :

$$C_0 + C_1(x - a) + C_2(x - a)^2 + \cdots + C_n(x - a)^n + \cdots = \sum_{n=0}^{\infty} C_n(x - a)^n.$$

We think of  $a$  as a constant. For any fixed  $x$ , the power series  $\sum C_n(x - a)^n$  is a series of numbers like those considered in Section 9.3. To investigate the convergence of a power series, we consider the partial sums, which in this case are the polynomials  $S_n(x) = C_0 + C_1(x - a) + C_2(x - a)^2 + \cdots + C_n(x - a)^n$ . As before, we consider the sequence<sup>7</sup>

$$S_0(x), S_1(x), S_2(x), \dots, S_n(x), \dots$$

For a fixed value of  $x$ , if this sequence of partial sums converges to a limit  $L$ , that is, if  $\lim_{n \rightarrow \infty} S_n(x) = L$ , then we say that the power series **converges** to  $L$  for this value of  $x$ .

A power series may converge for some values of  $x$  and not for others.

**Example 1** Determine whether the power series  $\sum_{n=0}^{\infty} \frac{x^n}{2^n}$  converges or diverges for

(a)  $x = -1$  (b)  $x = 3$

**Solution** (a) Substituting  $x = -1$ , we have

$$\sum_{n=0}^{\infty} \frac{x^n}{2^n} = \sum_{n=0}^{\infty} \frac{(-1)^n}{2^n} = \sum_{n=0}^{\infty} \left(-\frac{1}{2}\right)^n.$$

This is a geometric series with ratio  $-1/2$ , so the series converges to  $1/(1 - (-1/2)) = 2/3$ .

<sup>7</sup>Here we have chosen to call the first term in the sequence  $S_0(x)$  rather than  $S_1(x)$  to correspond to the power of  $(x - a)$ .

(b) Substituting  $x = 3$ , we have

$$\sum_{n=0}^{\infty} \frac{x^n}{2^n} = \sum_{n=0}^{\infty} \frac{3^n}{2^n} = \sum_{n=0}^{\infty} \left(\frac{3}{2}\right)^n.$$

This is a geometric series with ratio greater than 1, so it diverges.

**Example 2** Find an expression for the general term of the series and use it to write the series using  $\sum$  notation:

$$\frac{(x-2)^4}{4} - \frac{(x-2)^6}{9} + \frac{(x-2)^8}{16} - \frac{(x-2)^{10}}{25} + \dots$$

**Solution** The series is about  $x = 2$  and the odd terms are zero. We use  $(x-2)^{2n}$  and begin with  $n = 2$ . Since the series alternates and is positive for  $n = 2$ , we multiply by  $(-1)^n$ . For  $n = 2$ , we divide by 4, for  $n = 3$  we divide by 9, and in general, we divide by  $n^2$ . One way to write this series is

$$\sum_{n=2}^{\infty} \frac{(-1)^n (x-2)^{2n}}{n^2}.$$

## Numerical and Graphical View of Convergence

Consider the series


$$(x-1) - \frac{(x-1)^2}{2} + \frac{(x-1)^3}{3} - \frac{(x-1)^4}{4} + \dots + (-1)^{n-1} \frac{(x-1)^n}{n} + \dots$$

To investigate the convergence of this series, we look at the sequence of partial sums graphed in Figure 9.10. The graph suggests that the partial sums converge for  $x$  in the interval  $(0, 2)$ . In Examples 4 and 6, we show that the series converges for  $0 < x \leq 2$ . This is called the *interval of convergence* of this series.

At  $x = 1.4$ , which is inside the interval, the series appears to converge quite rapidly:

$$\begin{aligned} S_5(1.4) &= 0.33698\dots & S_7(1.4) &= 0.33653\dots \\ S_6(1.4) &= 0.33630\dots & S_8(1.4) &= 0.33645\dots \end{aligned}$$

Table 9.1 shows the results of using  $x = 1.9$  and  $x = 2.3$  in the power series. For  $x = 1.9$ , which is inside the interval of convergence but close to an endpoint, the series converges, though rather slowly. For  $x = 2.3$ , which is outside the interval of convergence, the series diverges: the larger the value of  $n$ , the more wildly the series oscillates. In fact, the contribution of the twenty-fifth term is about 28; the contribution of the hundredth term is about  $-2,500,000,000$ . Figure 9.10 shows the interval of convergence and the partial sums.



**Figure 9.10:** Partial sums for series in Example 4 converge for  $0 < x < 2$

**Table 9.1** Partial sums for series in Example 4 with  $x = 1.9$  inside interval of convergence and  $x = 2.3$  outside


$n$	$S_n(1.9)$	$n$	$S_n(2.3)$
2	0.495	2	0.455
5	0.69207	5	1.21589
8	0.61802	8	0.28817
11	0.65473	11	1.71710
14	0.63440	14	-0.70701

Notice that the interval of convergence,  $0 < x \leq 2$ , is centered on  $x = 1$ . Since the interval extends one unit on either side, we say the *radius of convergence* of this series is 1.

## Intervals of Convergence

Each power series falls into one of the three following cases, characterized by its *radius of convergence*,  $R$ .

- The series converges only for  $x = a$ ; the **radius of convergence** is defined to be  $R = 0$ .
- The series converges for all values of  $x$ ; the **radius of convergence** is defined to be  $R = \infty$ .
- There is a positive number  $R$ , called the **radius of convergence**, such that the series converges for  $|x - a| < R$  and diverges for  $|x - a| > R$ . See Figure 9.11.
- The **interval of convergence** is the interval between  $a - R$  and  $a + R$ , including any endpoint where the series converges.



**Figure 9.11:** Radius of convergence,  $R$ , determines an interval, centered at  $x = a$ , in which the series converges

There are some series whose radius of convergence we already know. For example, the geometric series

$$1 + x + x^2 + \cdots + x^n + \cdots$$

converges for  $|x| < 1$  and diverges for  $|x| \geq 1$ , so its radius of convergence is 1. Similarly, the series

$$1 + \frac{x}{3} + \left(\frac{x}{3}\right)^2 + \cdots + \left(\frac{x}{3}\right)^n + \cdots$$

converges for  $|x/3| < 1$  and diverges for  $|x/3| \geq 1$ , so its radius of convergence is 3.

The next theorem gives a method of computing the radius of convergence for many series. To find the values of  $x$  for which the power series  $\sum_{n=0}^{\infty} C_n(x - a)^n$  converges, we use the ratio test. Writing  $a_n = C_n(x - a)^n$  and assuming  $C_n \neq 0$  and  $x \neq a$ , we have

$$\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = \lim_{n \rightarrow \infty} \frac{|C_{n+1}(x - a)^{n+1}|}{|C_n(x - a)^n|} = \lim_{n \rightarrow \infty} \frac{|C_{n+1}||x - a|}{|C_n|} = |x - a| \lim_{n \rightarrow \infty} \frac{|C_{n+1}|}{|C_n|}.$$

**Case 1.** Suppose  $\lim_{n \rightarrow \infty} |a_{n+1}|/|a_n|$  is infinite. Then the ratio test shows that the power series converges only for  $x = a$ . The radius of convergence is  $R = 0$ .

**Case 2.** Suppose  $\lim_{n \rightarrow \infty} |a_{n+1}|/|a_n| = 0$ . Then the ratio test shows that the power series converges for all  $x$ . The radius of convergence is  $R = \infty$ .

**Case 3.** Suppose  $\lim_{n \rightarrow \infty} |a_{n+1}|/|a_n| = K|x - a|$ , where  $\lim_{n \rightarrow \infty} |C_{n+1}|/|C_n| = K$ . In Case 1,  $K$  does not exist; in Case 2,  $K = 0$ . Thus, we can assume  $K$  exists and  $K \neq 0$ , and we can define  $R = 1/K$ . Then we have

$$\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = K|x - a| = \frac{|x - a|}{R},$$

so the ratio test tells us that the power series:

- Converges for  $\frac{|x-a|}{R} < 1$ ; that is, for  $|x-a| < R$
- Diverges for  $\frac{|x-a|}{R} > 1$ ; that is, for  $|x-a| > R$ .

The results are summarized in the following theorem.

### Theorem 9.10: Method for Computing Radius of Convergence

To calculate the radius of convergence,  $R$ , for the power series  $\sum_{n=0}^{\infty} C_n(x-a)^n$ , use the ratio test with  $a_n = C_n(x-a)^n$ .

- If  $\lim_{n \rightarrow \infty} |a_{n+1}|/|a_n|$  is infinite, then  $R = 0$ .
- If  $\lim_{n \rightarrow \infty} |a_{n+1}|/|a_n| = 0$ , then  $R = \infty$ .
- If  $\lim_{n \rightarrow \infty} |a_{n+1}|/|a_n| = K|x-a|$ , where  $K$  is finite and nonzero, then  $R = 1/K$ .

Note that the ratio test does not tell us anything if  $\lim_{n \rightarrow \infty} |a_{n+1}|/|a_n|$  fails to exist, which can occur, for example, if some of the  $C_n$ s are zero.

A proof that a power series has a radius of convergence and of Theorem 9.10 is given in the online theory supplement. To understand these facts informally, we can think of a power series as being like a geometric series whose coefficients vary from term to term. The radius of convergence depends on the behavior of the coefficients: if there are constants  $C$  and  $K$  such that for larger and larger  $n$ ,

$$|C_n| \approx CK^n,$$

then it is plausible that  $\sum C_n x^n$  and  $\sum CK^n x^n = \sum C(Kx)^n$  converge or diverge together. The geometric series  $\sum C(Kx)^n$  converges for  $|Kx| < 1$ , that is, for  $|x| < 1/K$ . We can find  $K$  using the ratio test, because

$$\frac{|a_{n+1}|}{|a_n|} = \frac{|C_{n+1}||x-a|^{n+1}|}{|C_n||x-a|^n|} \approx \frac{CK^{n+1}|x-a|^{n+1}|}{CK^n|x-a|^n|} = K|x-a|.$$

**Example 3** Show that the following power series converges for all  $x$ :

$$1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!} + \cdots.$$

**Solution** Because  $C_n = 1/n!$ , none of the  $C_n$ s are zero and we can use the ratio test:

$$\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = |x| \lim_{n \rightarrow \infty} \frac{|C_{n+1}|}{|C_n|} = |x| \lim_{n \rightarrow \infty} \frac{1/(n+1)!}{1/n!} = |x| \lim_{n \rightarrow \infty} \frac{n!}{(n+1)!} = |x| \lim_{n \rightarrow \infty} \frac{1}{n+1} = 0.$$

This gives  $R = \infty$ , so the series converges for all  $x$ . We see in Chapter 10 that it converges to  $e^x$ .

**Example 4** Determine the radius of convergence of the series

$$(x-1) - \frac{(x-1)^2}{2} + \frac{(x-1)^3}{3} - \frac{(x-1)^4}{4} + \cdots + (-1)^{n-1} \frac{(x-1)^n}{n} + \cdots.$$

What does this tell us about the interval of convergence of this series?



**Solution** The general term of the series is  $(x-1)^n/n$  if  $n$  is odd and  $-(x-1)^n/n$  if  $n$  is even, so  $C_n = (-1)^{n-1}/n$ , and we can use the ratio test. We have

$$\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = |x-1| \lim_{n \rightarrow \infty} \frac{|C_{n+1}|}{|C_n|} = |x-1| \lim_{n \rightarrow \infty} \frac{|(-1)^n/(n+1)|}{|(-1)^{n-1}/n|} = |x-1| \lim_{n \rightarrow \infty} \frac{n}{n+1} = |x-1|.$$

Thus,  $K = 1$  in Theorem 9.10, so the radius of convergence is  $R = 1$ . The power series converges for  $|x-1| < 1$  and diverges for  $|x-1| > 1$ , so the series converges for  $0 < x < 2$ . Notice that the radius of convergence does not tell us what happens at the endpoints,  $x = 0$  and  $x = 2$ . We see in Chapter 10 that the series converges to  $\ln x$  for  $0 < x \leq 2$ .

The ratio test requires  $\lim_{n \rightarrow \infty} |a_{n+1}|/|a_n|$  to exist for  $a_n = C_n(x-a)^n$ . What happens if some of the coefficients  $C_n$  are zero? Then we use the fact that an infinite series can be written in several ways and pick one in which the terms are nonzero. For example, we think of the power series

$$x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots + (-1)^{n-1} \frac{x^{2n-1}}{(2n-1)!} + \cdots$$

as the series with  $a_1 = x$  and  $a_2 = -x^3/3!, \dots$ , so  $a_n = (-1)^{n-1}x^{2n-1}/(2n-1)!$ . With this choice of  $a_n$ , all  $a_n \neq 0$ , so we can use the ratio test.<sup>8</sup>

**Example 5** Find the radius and interval of convergence of the series

$$x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots + (-1)^{n-1} \frac{x^{2n-1}}{(2n-1)!} + \cdots$$

**Solution** We take

$$a_n = (-1)^{n-1} \frac{x^{2n-1}}{(2n-1)!},$$

so that, replacing  $n$  by  $n+1$ , we have

$$a_{n+1} = (-1)^{(n+1)-1} \frac{x^{2(n+1)-1}}{(2(n+1)-1)!} = (-1)^n \frac{x^{2n+1}}{(2n+1)!}.$$

Thus,

$$\frac{|a_{n+1}|}{|a_n|} = \frac{\left| (-1)^n \frac{x^{2n+1}}{(2n+1)!} \right|}{\left| (-1)^{n-1} \frac{x^{2n-1}}{(2n-1)!} \right|} = \left| \frac{(-1)^n x^{2n+1} (2n-1)!}{(-1)^{n-1} x^{2n-1} (2n+1)!} \right| = \left| \frac{(-1) x^2}{(2n+1) 2n} \right| = \frac{x^2}{(2n+1) 2n}.$$

Because

$$\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = \lim_{n \rightarrow \infty} \frac{x^2}{(2n+1) 2n} = 0,$$

we have  $\lim_{n \rightarrow \infty} |a_{n+1}|/|a_n| = 0 < 1$  for all  $x$ . Thus, the ratio test guarantees that the power series converges for every  $x$ . The radius of convergence is infinite and the interval of convergence is all  $x$ . We see in Chapter 10 that the series converges to  $\sin x$ .

<sup>8</sup>We do not take  $a_1 = x, a_2 = 0, a_3 = -x^3/3!, a_4 = 0, \dots$  because then  $\lim_{n \rightarrow \infty} |a_{n+1}|/|a_n|$  does not exist.

## What Happens at the Endpoints of the Interval of Convergence?

The ratio test does not tell us whether a power series converges at the endpoints of its interval of convergence,  $x = a \pm R$ . There is no simple theorem that answers this question. Since substituting  $x = a \pm R$  converts the power series to a series of numbers, the tests in Sections 9.3 and 9.4 are often useful. See Examples 6 and 7.

**Example 6** Determine the interval of convergence of the series

$$(x-1) - \frac{(x-1)^2}{2} + \frac{(x-1)^3}{3} - \frac{(x-1)^4}{4} + \cdots + (-1)^{n-1} \frac{(x-1)^n}{n} + \cdots.$$

**Solution** In Example 4 on page 493, we showed that this series has  $R = 1$ ; it converges for  $0 < x < 2$  and diverges for  $x < 0$  or  $x > 2$ . We need to determine whether it converges at the endpoints of the interval of convergence,  $x = 0$  and  $x = 2$ . At  $x = 2$ , we have the series

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots + \frac{(-1)^{n-1}}{n} + \cdots.$$

This is an alternating series with  $a_n = 1/(n+1)$ , so by the alternating series test (Theorem 9.8), it converges. At  $x = 0$ , we have the series

$$-1 - \frac{1}{2} - \frac{1}{3} - \frac{1}{4} - \cdots - \frac{1}{n} - \cdots.$$

This is the negative of the harmonic series, so it diverges. Therefore, the interval of convergence is  $0 < x \leq 2$ . The right endpoint is included and the left endpoint is not.

**Example 7** Find the radius and interval of convergence of the series

$$1 + 2^2x^2 + 2^4x^4 + 2^6x^6 + \cdots + 2^{2n}x^{2n} + \cdots.$$

**Solution** If we take  $a_n = 2^n x^n$  for  $n$  even and  $a_n = 0$  for  $n$  odd,  $\lim_{n \rightarrow \infty} |a_{n+1}|/|a_n|$  does not exist. Therefore, for this series we take

$$a_n = 2^{2n} x^{2n},$$

so that, replacing  $n$  by  $n+1$ , we have

$$a_{n+1} = 2^{2(n+1)} x^{2(n+1)} = 2^{2n+2} x^{2n+2}.$$

Thus,

$$\frac{|a_{n+1}|}{|a_n|} = \frac{|2^{2n+2} x^{2n+2}|}{|2^{2n} x^{2n}|} = |2^2 x^2| = 4x^2.$$

We have

$$\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = 4x^2.$$

The ratio test guarantees that the power series converges if  $4x^2 < 1$ , that is, if  $|x| < \frac{1}{2}$ . The radius of convergence is  $\frac{1}{2}$ . The series converges for  $-\frac{1}{2} < x < \frac{1}{2}$  and diverges for  $x < -\frac{1}{2}$  or  $x > \frac{1}{2}$ . At  $x = \pm \frac{1}{2}$ , all the terms in the series are 1, so the series diverges (by Theorem 9.2 Property 3). Thus, the interval of convergence is  $-\frac{1}{2} < x < \frac{1}{2}$ .

## Exercises and Problems for Section 9.5

## Exercises

Which of the series in Exercises 1–4 are power series?

- $x - x^3 + x^6 - x^{10} + x^{15} - \dots$
- $\frac{1}{x} + \frac{1}{x^2} + \frac{1}{x^3} + \frac{1}{x^4} + \dots$
- $1 + x + (x-1)^2 + (x-2)^3 + (x-3)^4 + \dots$
- $x^7 + x + 2$

In Exercises 5–10, find an expression for the general term of the series. Give the starting value of the index ( $n$  or  $k$  for example).

- $\frac{1}{2}x + \frac{1 \cdot 3}{2^2 \cdot 2!}x^2 + \frac{1 \cdot 3 \cdot 5}{2^3 \cdot 3!}x^3 + \dots$
- $px + \frac{p(p-1)}{2!}x^2 + \frac{p(p-1)(p-2)}{3!}x^3 + \dots$
- $1 - \frac{(x-1)^2}{2!} + \frac{(x-1)^4}{4!} - \frac{(x-1)^6}{6!} + \dots$
- $(x-1)^3 - \frac{(x-1)^5}{2!} + \frac{(x-1)^7}{4!} - \frac{(x-1)^9}{6!} + \dots$
- $\frac{x-a}{1} + \frac{(x-a)^2}{2 \cdot 2!} + \frac{(x-a)^3}{4 \cdot 3!} + \frac{(x-a)^4}{8 \cdot 4!} + \dots$
- $2(x+5)^3 + 3(x+5)^5 + \frac{4(x+5)^7}{2!} + \frac{5(x+5)^9}{3!} + \dots$

Use the ratio test to find the radius of convergence of the power series in Exercises 11–21.

- $\sum_{n=0}^{\infty} (5x)^n$
- $\sum_{n=0}^{\infty} n^3 x^n$
- $\sum_{n=0}^{\infty} \frac{(n+1)x^n}{2^n + n}$
- $\sum_{n=1}^{\infty} \frac{2^n (x-1)^n}{n}$
- $\sum_{n=1}^{\infty} \frac{(x-3)^n}{n2^n}$
- $\sum_{n=1}^{\infty} \frac{x^{2n}}{(2n)!}$
- $x - \frac{x^2}{4} + \frac{x^3}{9} - \frac{x^4}{16} + \frac{x^5}{25} - \dots$
- $1 + 2x + \frac{4x^2}{2!} + \frac{8x^3}{3!} + \frac{16x^4}{4!} + \frac{32x^5}{5!} + \dots$
- $1 + 2x + \frac{4!x^2}{(2!)^2} + \frac{6!x^3}{(3!)^2} + \frac{8!x^4}{(4!)^2} + \frac{10!x^5}{(5!)^2} + \dots$
- $3x + \frac{5}{2}x^2 + \frac{7}{3}x^3 + \frac{9}{4}x^4 + \frac{11}{5}x^5 + \dots$
- $x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots$

## Problems

22. (a) Determine the radius of convergence of the series

$$x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots + (-1)^{n-1} \frac{x^n}{n} + \dots$$

What does this tell us about the interval of convergence of this series?

- (b) Investigate convergence at the end points of the interval of convergence of this series.

23. Show that the series  $\sum_{n=1}^{\infty} \frac{(2x)^n}{n}$  converges for  $|x| < 1/2$ .

Investigate whether the series converges for  $x = 1/2$  and  $x = -1/2$ .

In Problems 24–29, find the interval of convergence.

- $\sum_{n=0}^{\infty} \frac{x^n}{3^n}$
- $\sum_{n=2}^{\infty} \frac{(x-3)^n}{n}$
- $\sum_{n=1}^{\infty} \frac{n^2 x^{2n}}{2^{2n}}$
- $\sum_{n=1}^{\infty} \frac{(-1)^n (x-5)^n}{2^n n^2}$
- $\sum_{n=1}^{\infty} \frac{x^n}{n!}$
- $\sum_{n=0}^{\infty} n! x^n$

In Problems 30–33, use the formula for the sum of a geometric series to find a power series centered at the origin that converges to the expression. For what values does the series converge?

- $\frac{1}{1+2z}$
- $\frac{2}{1+y^2}$
- $\frac{3}{1-z/2}$
- $\frac{8}{4+y}$

34. For constant  $p$ , find the radius of convergence of the binomial power series:<sup>9</sup>

$$1 + px + \frac{p(p-1)x^2}{2!} + \frac{p(p-1)(p-2)x^3}{3!} + \dots$$

- Show that if  $C_0 + C_1x + C_2x^2 + C_3x^3 + \dots$  converges for  $|x| < R$  with  $R$  given by the ratio test, then so does  $C_1 + 2C_2x + 3C_3x^2 + \dots$ .
- The series  $\sum C_n x^n$  converges at  $x = -5$  and diverges at  $x = 7$ . What can you say about its radius of convergence?

<sup>9</sup>For an explanation of the name, see Section 10.2.

37. The series  $\sum C_n(x+7)^n$  converges at  $x = 0$  and diverges at  $x = -17$ . What can you say about its radius of convergence?
38. Suppose that the power series  $\sum C_n x^n$  converges when  $x = -4$  and diverges when  $x = 7$ . Which of the following are true, false or not possible to determine?
- The power series converges when  $x = 10$ .
  - The power series converges when  $x = 3$ .
  - The power series diverges when  $x = 1$ .
  - The power series diverges when  $x = 6$ .
39. If  $\sum C_n(x-3)^n$  converges at  $x = 7$  and diverges at  $x = 10$ , what can you say about the convergence at  $x = 11$ ? At  $x = 5$ ? At  $x = 0$ ?

40. Bessel functions are important in such diverse areas as describing planetary motion and the shape of a vibrating drumhead. The Bessel function of order 0 is defined by

$$J(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{2^{2n} (n!)^2}.$$

- Find the domain of  $J(x)$  by finding the interval of convergence for this power series.
  - Find  $J(0)$ .
  - Find the partial sum polynomials  $S_0, S_1, S_2, S_3, S_4$ .
  - Estimate  $J(1)$  to three decimal places.
  - Use your answer to part (d) to estimate  $J(-1)$ .
41. For all  $x$ -values for which it converges, the function  $f$  is defined by the series

$$f(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!}.$$

- What is  $f(0)$ ?
- What is the domain of  $f$ ?
- Assuming that  $f'$  can be calculated by differentiating the series term-by-term, find the series for  $f'(x)$ . What do you notice?
- Guess what well-known function  $f$  is.

42. From Example 5 on page 494, we know the following series converges for all  $x$ . We define  $g(x)$  to be its sum:

$$g(x) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n-1}}{(2n-1)!}.$$

- Is  $g(x)$  odd, even, or neither? What is  $g(0)$ ?
- Assuming that derivatives can be computed term-by-term, show that  $g''(x) = -g(x)$ .
- Guess what well-known function  $g$  might be. Check your guess using  $g(0)$  and  $g'(0)$ .

43. The functions  $p(x)$  and  $q(x)$  are defined by the series

$$p(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}, \quad q(x) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n-1}}{(2n-1)!}.$$

Assuming that these series converge for all  $x$  and that multiplication can be done term-by-term:

- Find the series for  $(p(x))^2 + (q(x))^2$  up to the term in  $x^6$ .
- Guess what well-known functions  $p$  and  $q$  could be.

## CHAPTER SUMMARY (see also Ready Reference at the end of the book)

- Sequences**  
Recursive definition, monotone, bounded, convergence.
- Geometric series**  
Finite sum, infinite sum.
- Harmonic series**
- Alternating series**
- Tests for convergence of series**  
Integral test,  $p$ -series, comparison test, limit comparison test, ratio test, alternating series test.  
Absolute and conditional convergence.
- Power series**  
Ratio test for radius of convergence, interval of convergence.

## REVIEW EXERCISES AND PROBLEMS FOR CHAPTER NINE

### Exercises

- Find the sum of the series  $b^5 + b^6 + b^7 + b^8 + b^9 + b^{10}$ .
- Find the sum of the series  $(0.5)^3 + (0.5)^4 + \cdots + (0.5)^k$ .
- Find the sum:  $\sum_{n=0}^{\infty} \frac{3^n + 5}{4^n}$

Do the sequences in Exercises 4–7 converge or diverge? If a sequence converges, find its limit.

- $\frac{3+4n}{5+7n}$
- $(-1)^n \frac{(n+1)}{n}$
- $\sin\left(\frac{\pi n}{4}\right)$
- $\frac{1}{n} + \ln n$

Use the integral test to decide whether the series in Exercises 8–11 converge or diverge.

$$8. \sum_{n=1}^{\infty} \frac{1}{(n+2)^2}$$

$$9. \sum_{n=1}^{\infty} \frac{3n^2 + 2n}{n^3 + n^2 + 1}$$

$$10. \sum_{n=0}^{\infty} ne^{-n^2}$$

$$11. \sum_{n=2}^{\infty} \frac{2}{n^2 - 1}$$

Use the ratio test to decide if the series in Exercises 12–15 converge or diverge.

$$12. \sum_{n=1}^{\infty} \frac{n}{2^n}$$

$$13. \sum_{n=1}^{\infty} \frac{1}{2^n n!}$$

$$14. \sum_{n=1}^{\infty} \frac{(n-1)!}{5^n}$$

$$15. \sum_{n=1}^{\infty} \frac{n!(n+1)!}{(2n)!}$$

Use the alternating series test to decide which of the series in Exercises 16–17 converge.

$$16. \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2 + 1}$$

$$17. \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{\sqrt{n^2 + 1}}$$

In Exercises 18–21 determine whether the series are absolutely convergent, conditionally convergent, or divergent.

$$18. \sum \frac{(-1)^n}{n^{1/2}}$$

$$19. \sum (-1)^n \frac{n}{n+1}$$

$$20. \sum \frac{(-1)^{n-1} \ln n}{n}$$

$$21. \sum \frac{(-1)^{n-1}}{\arctan n}$$

Use the comparison test to confirm the statements in Exercises 22–23.

$$22. \sum_{n=1}^{\infty} \left(\frac{1}{3}\right)^n \text{ converges, so } \sum_{n=1}^{\infty} \left(\frac{n^2}{3n^2 + 4}\right)^n \text{ converges.}$$

$$23. \sum_{n=1}^{\infty} \frac{1}{n} \text{ diverges, so } \sum_{n=1}^{\infty} \frac{1}{n \sin^2 n} \text{ diverges.}$$

In Exercises 24–27 use the limit comparison test to determine whether the series converges or diverges.

$$24. \sum \frac{\sqrt{n-1}}{n^2 + 3}$$

$$25. \sum \frac{n^3 - 2n^2 + n + 1}{n^5 - 2}$$

$$26. \sum \sin \frac{1}{n^2}$$

$$27. \sum \frac{1}{\sqrt{n^3 - 1}}$$

Determine which of the series in Exercises 28–47 converge.

$$28. \sum_{n=1}^{\infty} \frac{1}{n+1}$$

$$29. \sum_{n=1}^{\infty} \frac{1}{n^3}$$

$$30. \sum_{n=3}^{\infty} \frac{2}{\sqrt{n-2}}$$

$$31. \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{\sqrt{n+1}}$$

$$32. \sum_{n=1}^{\infty} \frac{n^2}{n^2 + 1}$$

$$33. \sum_{n=1}^{\infty} \frac{n^2}{n^3 + 1}$$

$$34. \sum_{n=1}^{\infty} \frac{3^n}{(2n)!}$$

$$35. \sum_{n=1}^{\infty} \frac{(2n)!}{(n!)^2}$$

$$36. \sum_{n=1}^{\infty} \frac{n^2 + 2^n}{n^2 2^n}$$

$$37. \sum_{n=1}^{\infty} 2^{-n} \frac{(n+1)}{(n+2)}$$

$$38. \sum_{n=1}^{\infty} (-1)^n \frac{2^n}{(2n+1)!}$$

$$39. \sum_{n=1}^{\infty} (-1)^n \frac{n+1}{\sqrt{n}}$$

$$40. \sum_{n=0}^{\infty} \frac{2 + 3^n}{5^n}$$

$$41. \sum_{n=1}^{\infty} \frac{1}{2 + \sin n}$$

$$42. \sum_{n=3}^{\infty} \frac{1}{(2n-5)^3}$$

$$43. \sum_{n=2}^{\infty} \frac{1}{n^3 - 3}$$

$$44. \sum_{n=1}^{\infty} \frac{\sin(n\pi/2)}{n^3}$$

$$45. \sum_{k=1}^{\infty} \ln \left(1 + \frac{1}{k}\right)$$

$$46. \sum_{n=1}^{\infty} \frac{n}{2^n}$$

$$47. \sum_{n=2}^{\infty} \frac{1}{(\ln n)^2}$$

In Exercises 48–53, find the radius of convergence.

$$48. \sum_{n=0}^{\infty} nx^n$$

$$49. \sum_{n=1}^{\infty} \frac{(2n)!x^n}{(n!)^2}$$

$$50. \sum_{n=0}^{\infty} (2^n + n^2)x^n$$

$$51. \sum_{n=0}^{\infty} \frac{x^n}{n! + 1}$$

$$52. x + 4x^2 + 9x^3 + 16x^4 + 25x^5 + \cdots$$

$$53. \frac{x}{3} + \frac{2x^2}{5} + \frac{3x^3}{7} + \frac{4x^4}{9} + \frac{5x^5}{11} + \cdots$$

In Exercises 54–57, find the interval of convergence.

$$54. \sum_{n=1}^{\infty} \frac{x^n}{3^n n^2}$$

$$55. \sum_{n=0}^{\infty} \frac{(-1)^n (x-2)^n}{5^n}$$

$$56. \sum_{n=1}^{\infty} \frac{(-1)^n x^n}{n}$$

$$57. \sum_{n=1}^{\infty} \frac{x^{2n+1}}{n!}$$

## Problems

58. For  $r > 0$ , how does the convergence of the following series depend on  $r$ ?

$$\sum_{n=1}^{\infty} \frac{n^r + r^n}{n^r r^n}$$

59. Suppose that  $\sum_{n=0}^{\infty} C_n(x-2)^n$  converges when  $x = 4$  and diverges when  $x = 6$ . Which of the following are true, false or not possible to determine? Give reasons for your answers.

- (a) The power series converges when  $x = 7$ .
- (b) The power series diverges when  $x = 1$ .
- (c) The power series converges when  $x = 0.5$ .
- (d) The power series diverges when  $x = 5$ .
- (e) The power series converges when  $x = -3$ .

60. For all the  $t$ -values for which it converges, the function  $h$  is defined by the series

$$h(t) = \sum_{n=0}^{\infty} (-1)^n \frac{t^{2n}}{(2n)!}.$$

- (a) What is the domain of  $h$ ?
- (b) Is  $h$  odd, even, or neither?
- (c) Assuming that derivatives can be computed term-by-term, show that

$$h''(t) = -h(t).$$

61. Peter wishes to create a retirement fund from which he can draw \$20,000 when he retires and the same amount at each anniversary of his retirement for 10 years. He plans to retire 20 years from now. What investment need he make today if he can get a return of 5% per year, compounded annually?
62. The extraction rate of a mineral is currently 12 million tons a year, but this rate is expected to fall by 5% each year. What minimum level of world reserves would allow extraction to continue indefinitely?
63. A new car costs \$30,000; it loses 10% of its value each year. Maintenance is \$500 the first year and increases by 20% annually.
- (a) Find a formula for  $l_n$ , the value lost by the car in year  $n$ .
  - (b) Find a formula for  $m_n$ , the maintenance expenses in year  $n$ .
  - (c) In what year do maintenance expenses first exceed the value lost by the car?

Problems 64–67 are about *bonds*, which are issued by a government to raise money. An individual who buys a \$1000 bond gives the government \$1000 and in return receives a fixed sum of money, called the *coupon*, every six months or every year for the life of the bond. At the time of the last coupon, the individual also gets the \$1000, or *principal* back.

64. What is the present value of a \$1000 bond which pays \$50 a year for 10 years, starting one year from now? Assume the interest rate is 6% per year, compounded annually.

65. What is the present value of a \$1000 bond which pays \$50 a year for 10 years, starting one year from now? Assume the interest rate is 4% per year, compounded annually.

66. (a) What is the present value of a \$1000 bond which pays \$50 a year for 10 years, starting one year from now? Assume the interest rate is 5% per year, compounded annually.
- (b) Since \$50 is 5% of \$1000, this bond is often called a 5% bond. What does your answer to part (a) tell you about the relationship between the principal and the present value of this bond when the interest rate is 5%?
- (c) If the interest rate is more than 5% per year, compounded annually, which is larger: the principal or the present value of the bond? Why do you think the bond is then described as *trading at a discount*?
- (d) If the interest rate is less than 5% per year, compounded annually, why is the bond described as *trading at a premium*?

67. In the nineteenth century, the railroads issued 100-year bonds. Consider a \$100 bond which paid \$5 a year, starting a year after it was sold. Assume interest rates are 4% per year, compounded annually.

- (a) What was such a bond worth on the day it was sold?
- (b) Suppose that instead of maturing in 100 years, the bond was to have paid \$5 a year forever. This time the principal, \$100, is never repaid. How much would such a bond be worth on the day of its sale?

68. Cephalexin is an antibiotic with a half-life in the body of 0.9 hours, taken in tablets of 250 mg every six hours.

- (a) What percentage of the cephalexin in the body at the start of a six-hour period is still there at the end (assuming no tablets are taken during that time)?
- (b) Write an expression for  $Q_1, Q_2, Q_3, Q_4$ , where  $Q_n$  mg, is the amount of cephalexin in the body right after the  $n^{\text{th}}$  tablet is taken.
- (c) Express  $Q_3, Q_4$  in closed-form and evaluate them.
- (d) Write an expression for  $Q_n$  and put it in closed-form.
- (e) If the patient keeps taking the tablets, use your answer to part (d) to find the quantity of cephalexin in the body in the long run, right after taking a pill.

69. Before World War I, the British government issued what are called *consols*, which pay the owner or his heirs a fixed amount of money every year forever. (Cartoonists of the time described aristocrats living off such payments as “pickled in consols.”) What should a person expect to pay for consols which pay £10 a year forever? Assume the first payment is one year from the date of purchase and that interest remains 4% per year, compounded annually. (£ denotes pounds, the British unit of currency.)
70. This problem illustrates how banks create credit and can thereby lend out more money than has been deposited. Suppose that initially \$100 is deposited in a bank. Experience has shown bankers that on average only 8% of the money deposited is withdrawn by the owner at any time. Consequently, bankers feel free to lend out 92% of their deposits. Thus \$92 of the original \$100 is loaned out to other customers (to start a business, for example). This \$92 will become someone else’s income and, sooner or later, will be redeposited in the bank. Then 92% of \$92, or  $\$92(0.92) = \$84.64$ , is loaned out again and eventually redeposited. Of the \$84.64, the bank again loans out 92%, and so on.
- (a) Find the total amount of money deposited in the bank as a result of these transactions.
- (b) The total amount of money deposited divided by the original deposit is called the *credit multiplier*. Calculate the credit multiplier for this example and explain what this number tells us.
71. Baby formula can contain bacteria which double in number every half hour at room temperature and every 10 hours in the refrigerator.<sup>10</sup> Suppose there are  $B_0$  bacteria initially.
- (a) Write a formula for
- $R_n$ , the number of bacteria  $n$  hours later if the baby formula is kept at room temperature.
  - $F_n$ , the number of bacteria  $n$  hours later if the baby formula is kept in the refrigerator.
  - $Y_n$ , the ratio of the number of bacteria at room temperature to the number of bacteria in the refrigerator.
- (b) How many hours does it take before there are a million times as many bacteria in baby formula kept at room temperature as in baby formula kept in the refrigerator?
72. Estimate  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{(2n-1)!}$  to within 0.01 of the actual sum of the series.
73. Is it possible to construct a convergent alternating series  $\sum_{n=1}^{\infty} (-1)^{n-1} a_n$  for which  $0 < a_{n+1} < a_n$  but  $\lim_{n \rightarrow \infty} a_n \neq 0$ ?
74. Suppose that  $0 \leq b_n \leq 2^n$  for all  $n$ . Give two examples of series  $\sum b_n$  that satisfy this condition, one that diverges and one that converges.
75. Show that if  $\sum a_n$  converges and  $\sum b_n$  diverges, then  $\sum (a_n + b_n)$  diverges. [Hint: Assume that  $\sum (a_n + b_n)$  converges and consider  $\sum (a_n + b_n) - \sum a_n$ .]
- Suppose that  $\sum a_n$  converges with  $a_n > 0$  for all  $n$ . Decide if the series in Problems 76–80 converge, diverge or if there is not enough information provided.
76.  $\sum a_n/n$       77.  $\sum 1/a_n$       78.  $\sum na_n$
79.  $\sum (a_n + a_n/2)$       80.  $\sum a_n^2$
81. Does  $\sum_{n=1}^{\infty} \left(\frac{1}{n} + \frac{1}{n}\right)$  converge or diverge? Does  $\sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n}\right)$  converge or diverge? Is the statement “If  $\sum a_n$  and  $\sum b_n$  diverge, then  $\sum (a_n + b_n)$  may or may not diverge” true?
82. Estimate the sum of the first 100,000 terms of the harmonic series,
- $$\sum_{k=1}^{100,000} \frac{1}{k},$$
- to the closest integer. [Hint: Use left- and right-hand sums of the function  $f(x) = 1/x$  on the interval from 1 to 100,000 with  $\Delta x = 1$ .]
83. Although the harmonic series does not converge, the partial sums grow very, very slowly. Take a right-hand sum approximating the integral of  $f(x) = 1/x$  on the interval  $[1, n]$ , with  $\Delta x = 1$ , to show that
- $$\frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots + \frac{1}{n} < \ln n.$$
- If a computer could add a million terms of the harmonic series each second, estimate the sum after one year.
84. Is the following argument true or false? Give reasons for your answer.
- Consider the infinite series  $\sum_{n=2}^{\infty} \frac{1}{n(n-1)}$ . Since  $\frac{1}{n(n-1)} = \frac{1}{n-1} - \frac{1}{n}$  we can write this series as
- $$\sum_{n=2}^{\infty} \frac{1}{n-1} - \sum_{n=2}^{\infty} \frac{1}{n}.$$
- For the first series  $a_n = 1/(n-1)$ . Since  $n-1 < n$  we have  $1/(n-1) > 1/n$  and so this series diverges by comparison with the divergent harmonic series  $\sum_{n=2}^{\infty} \frac{1}{n}$ . The second series is the divergent harmonic series. Since both series diverge, their difference also diverges.

<sup>10</sup>Iverson, C. and Forsythe, F., reported in “Baby Food Could Trigger Meningitis,” [www.newscientist.com](http://www.newscientist.com), June 3, 2004.



## CHECK YOUR UNDERSTANDING

Decide if the statements in Problems 1–45 are true or false. Give an explanation for your answer.

- You can tell if a sequence converges by looking at the first 1000 terms.
- If the terms  $s_n$  of a convergent sequence are all positive then  $\lim_{n \rightarrow \infty} s_n$  is positive.
- If the sequence  $s_n$  of positive terms is unbounded, then the sequence has a term greater than a million.
- If the sequence  $s_n$  of positive terms is unbounded, then the sequence has an infinite number of terms greater than a million.
- If a sequence  $s_n$  is convergent, then the terms  $s_n$  tend to zero as  $n$  increases.
- If a series converges, then the sequence of partial sums of the series also converges.
- A monotone sequence can not have both positive and negative terms.
- If a monotone sequence of positive terms does not converge, then it has a term greater than a million.
- If the terms  $s_n$  of a sequence alternate in sign, then the sequence converges.
- If all terms  $s_n$  of a sequence are less than a million, then the sequence is bounded.
- $\sum_{n=1}^{\infty} (x - n)^n$  is a power series.
- If the power series  $\sum C_n x^n$  converges for  $x = 2$ , then it converges for  $x = 1$ .
- If the power series  $\sum C_n x^n$  converges for  $x = 1$ , then the power series converges for  $x = 2$ .
- If the power series  $\sum C_n x^n$  does not converge for  $x = 1$ , then the power series does not converge for  $x = 2$ .
- If  $0 \leq a_n \leq b_n$  and  $\sum a_n$  converges, then  $\sum b_n$  converges.
- If  $0 \leq a_n \leq b_n$  and  $\sum a_n$  diverges, then  $\sum b_n$  diverges.
- If  $b_n \leq a_n \leq 0$  and  $\sum b_n$  converges, then  $\sum a_n$  converges.
- If  $\sum a_n$  converges, then  $\sum |a_n|$  converges.
- If  $\sum |a_n + b_n|$  converges, then  $\sum |a_n|$  and  $\sum |b_n|$  converge.
- If  $\sum a_n$  converges, then  $\lim_{n \rightarrow \infty} |a_{n+1}|/|a_n| \neq 1$ .
- $\sum_{n=0}^{\infty} (-1)^n \cos(\pi n)$  is an alternating series.
- $\sum_{n=1}^{\infty} (1 + (-1)^n)$  is a series of nonnegative terms.
- The series  $\sum_{n=0}^{\infty} (-1)^n 2^n$  converges.
- The series  $\sum_{n=1}^{\infty} 2^{(-1)^n}$  converges.
- If  $\sum a_n$  converges, then  $\sum (-1)^n a_n$  converges.
- If an alternating series converges by the alternating series test, then the error in using the first  $n$  terms of the series to approximate the entire series is less in magnitude than the first term omitted.
- If an alternating series converges, then the error in using the first  $n$  terms of the series to approximate the entire series is less in magnitude than the first term omitted.
- If  $\sum |a_n|$  converges, then  $\sum (-1)^n |a_n|$  converges.
- To find the sum of the alternating harmonic series  $\sum (-1)^{n-1}/n$  to within 0.01 of the true value, we can sum the first 100 terms.
- $\sum C_n (x - 1)^n$  and  $\sum C_n x^n$  have the same radius of convergence.
- If  $\sum C_n x^n$  and  $\sum B_n x^n$  have the same radius of convergence, then the coefficients,  $C_n$  and  $B_n$ , must be equal.
- If a series  $\sum a_n$  converges, then the terms,  $a_n$ , tend to zero as  $n$  increases.
- If the terms,  $a_n$ , of a series tend to zero as  $n$  increases, then the series  $\sum a_n$  converges.
- If  $\sum a_n$  does not converge and  $\sum b_n$  does not converge, then  $\sum a_n b_n$  does not converge.
- If  $\sum a_n b_n$  converges, then  $\sum a_n$  and  $\sum b_n$  converge.
- If  $\sum a_n$  is absolutely convergent, then it is convergent.
- If  $\sum a_n$  is conditionally convergent, then it is absolutely convergent.
- If a power series converges at one endpoint of its interval of convergence, then it converges at the other.
- If  $a_n > 0.5b_n > 0$  for all  $n$  and  $\sum b_n$  diverges, then  $\sum a_n$  diverges.
- A power series always converges at at least one point.
- If the power series  $\sum C_n x^n$  converges at  $x = 10$ , then it converges at  $x = -9$ .
- If the power series  $\sum C_n x^n$  converges at  $x = 10$ , then it converges at  $x = -10$ .
- $-5 < x \leq 7$  is a possible interval of convergence of a power series.
- $-3 < x < 2$  could be the interval of convergence of  $\sum C_n x^n$ .
- If  $-11 < x < 1$  is the interval of convergence of  $\sum C_n (x - a)^n$ , then  $a = -5$ .

## PROJECTS FOR CHAPTER NINE

1. A Definition of  $e$ 

We show that the sequence  $s_n = \left(1 + \frac{1}{n}\right)^n$  converges; its limit can be used to define  $e$ .

- (a) For a fixed integer  $n > 0$ , let  $f(x) = (n+1)x^n - nx^{n+1}$ . For  $x > 1$ , show  $f$  is decreasing and that  $f(x) < 1$ . Hence, for  $x > 1$ ,

$$x^n(n+1-nx) < 1.$$

- (b) Substitute the following  $x$ -value into the inequality from part (a)

$$x = \frac{1 + 1/n}{1 + 1/(n+1)},$$

and show that

$$x^n \left( \frac{n+1}{n+2} \right) < 1.$$

- (c) Use the inequality from part (b) to show that  $s_n < s_{n+1}$  for all  $n > 0$ . Conclude the sequence is increasing.  
 (d) Substitute  $x = 1 + 1/2n$  into the inequality from part (a) to show that

$$\left(1 + \frac{1}{2n}\right)^n < 2.$$

- (e) Use the inequality from part (d) to show  $s_{2n} < 4$ . Conclude the sequence is bounded.  
 (f) Use parts (c) and (e) to show that the sequence has a limit.

## 2. Probability of Winning in Sports

In certain sports, winning a game requires a lead of two points. That is, if the score is tied you have to score two points in a row to win.

- (a) For some sports (e.g. tennis), a point is scored every play. Suppose your probability of scoring the next point is always  $p$ . Then, your opponent's probability of scoring the next point is always  $1 - p$ .
- What is the probability that you win the next two points?
  - What is the probability that you and your opponent split the next two points, that is, that neither of you wins both points?
  - What is the probability that you split the next two points but you win the two after that?
  - What is the probability that you either win the next two points or split the next two and then win the next two after that?
  - Give a formula for your probability  $w$  of winning a tied game.
  - Compute your probability of winning a tied game when  $p = 0.5$ ; when  $p = 0.6$ ; when  $p = 0.7$ ; when  $p = 0.4$ . Comment on your answers.
- (b) In other sports (e.g. volleyball), you can score a point only if it is your turn, with turns alternating until a point is scored. Suppose your probability of scoring a point when it is your turn is  $p$ , and your opponent's probability of scoring a point when it is her turn is  $q$ .
- Find a formula for the probability  $S$  that you are the first to score the next point, assuming it is currently your turn.
  - Suppose that if you score a point, the next turn is yours. Using your answers to part (a) and your formula for  $S$ , compute the probability of winning a tied game (if you need two points in a row to win).
    - Assume  $p = 0.5$  and  $q = 0.5$  and it is your turn.
    - Assume  $p = 0.6$  and  $q = 0.5$  and it is your turn.

**3. Prednisone**

Prednisone is often prescribed for acute asthma attacks. For 5 mg tablets, typical instructions are: “Take 8 tablets the first day, 7 the second, and decrease by one tablet each day until all tablets are gone.” Prednisone decays exponentially in the body, and 24 hours after taking  $k$  mg, there are  $kx$  mg in the body.

- (a) Write formulas involving  $x$  for the amount of prednisone in the body
  - (i) 24 hours after taking the first dose (of 8 tablets), right before taking the second dose (of 7 tablets).
  - (ii) Immediately after taking the second dose (of 7 tablets).
  - (iii) Immediately after taking the third dose (of 6 tablets).
  - (iv) Immediately after taking the eighth dose (of 1 tablet).
  - (v) 24 hours after taking the eighth dose.
  - (vi)  $n$  days after taking the eighth dose.
- (b) Find a closed form for the sum  $T = 8x^7 + 7x^6 + 6x^5 + \cdots + 2x + 1$ , which is the number of prednisone tablets in the body immediately after taking the eighth dose.
- (c) If a patient takes all the prednisone tablets as prescribed, how many days after taking the eighth dose is there less than 3% of a prednisone tablet in the patient’s body? The half-life of prednisone is about 24 hours.
- (d) A patient is prescribed  $n$  tablets of prednisone the first day,  $n - 1$  the second, and one tablet fewer each day until all tablets are gone. Write a formula that represents  $T_n$ , the number of prednisone tablets in the body immediately after taking all tablets. Find a closed form sum for  $T_n$ .

